



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 :

A61K 31/40, C07C 209/34

A1

(11) International Publication Number:

WO 96/40116

(43) International Publication Date: 19 December 1996 (19.12.96)

(21) International Application Number: PCT/US96/08903

(22) International Filing Date: 5 June 1996 (05.06.96)

(30) Priority Data:

485,323

7 June 1995 (07.06.95)

US

(71) Applicant: SUGEN, INC. [US/US]; 515 Galveston Drive,  
Redwood City, CA 94063 (US).(72) Inventors: TANG, Peng, Chor; 827 Camino Ricardo, Moraga,  
CA 94556 (US). SUN, Li; 64 Rockharbor Lane, Foster City,  
CA 94404 (US). MCMAHON, Gerald; 1800 Schultz Road,  
Kenwood, CA 95452 (US).(74) Agents: CORUZZI, Laura, A. et al.; Pennie & Edmonds, 1155  
Avenue of the Americas, New York, NY 10036 (US).(81) Designated States: AL, AM, AU, AZ, BB, BG, BR, BY, CA,  
CN, CZ, EE, FI, GE, HU, IL, IS, JP, KG, KP, KR, KZ,  
LK, LR, LS, LT, LV, MD, MG, MK, MN, MX, NO, NZ,  
PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, UZ, VN,  
ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: INDOLINONE COMPOUNDS FOR THE TREATMENT OF DISEASE

(57) Abstract

The present invention relates to organic molecules capable of modulating tyrosine kinase signal transduction in order to regulate, modulate and/or inhibit abnormal cell proliferation.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## INDOLINONE COMPOUNDS FOR THE TREATMENT OF DISEASE

This application is a continuation-in-part of U.S. Patent Application Serial No. 08/485,323, filed June 7, 1995, which is hereby incorporated by reference in its entirety.

5

## 1. INTRODUCTION

The present invention relates to novel compounds capable of modulating, regulating and/or inhibiting tyrosine kinase signal transduction. The present invention is also directed to methods of regulating, modulating or inhibiting tyrosine kinases, whether of the receptor or non-receptor class, for the prevention and/or treatment of disorders related to unregulated tyrosine kinase signal transduction, including cell proliferative and metabolic disorders.

15

## 2. BACKGROUND OF THE INVENTION

Protein tyrosine kinases (PTKs) comprise a large and diverse class of proteins having enzymatic activity. The PTKs play an important role in the control of cell growth and differentiation (for review, see Schlessinger & Ullrich, 1992, *Neuron* 9:383-391).

For example, receptor tyrosine kinase mediated signal transduction is initiated by extracellular interaction with a specific growth factor (ligand), followed by receptor dimerization, transient stimulation of the intrinsic protein tyrosine kinase activity and phosphorylation. Binding sites are thereby created for intracellular signal transduction molecules and lead to the formation of complexes with a spectrum of cytoplasmic signalling molecules that facilitate the appropriate cellular response (e.g., cell division, metabolic effects to the extracellular microenvironment). See, Schlessinger and Ullrich, 1992, *Neuron* 9:303-391.

With respect to receptor tyrosine kinases, it has been shown also that tyrosine phosphorylation sites function as high-affinity binding sites for SH2 (src homology) domains of signaling molecules. Fantl et al., 1992, *Cell* 69:413-423; Songyang et al., 1994, *Mol. Cell. Biol.* 14:2777-2785;

Songyang et al., 1993, *Cell* 72:767-778; and Koch et al., 1991, *Science* 252:668-678. Several intracellular substrate proteins that associate with receptor tyrosine kinases (RTKs) have been identified. They may be divided into two principal groups: (1) substrates which have a catalytic domain; and (2) substrates which lack such domain but serve as adapters and associate with catalytically active molecules. Songyang et al., 1993, *Cell* 72:767-778. The specificity of the interactions between receptors or proteins and SH2 domains of their substrates is determined by the amino acid residues immediately surrounding the phosphorylated tyrosine residue. Differences in the binding affinities between SH2 domains and the amino acid sequences surrounding the phosphotyrosine residues on particular receptors are consistent with the observed differences in their substrate phosphorylation profiles. Songyang et al., 1993, *Cell* 72:767-778. These observations suggest that the function of each receptor tyrosine kinase is determined not only by its pattern of expression and ligand availability but also by the array of downstream signal transduction pathways that are activated by a particular receptor. Thus, phosphorylation provides an important regulatory step which determines the selectivity of signaling pathways recruited by specific growth factor receptors, as well as differentiation factor receptors.

Aberrant expression or mutations in the PTKs have been shown to lead to either uncontrolled cell proliferation (e.g. malignant tumor growth) or to defects in key developmental processes. Consequently, the biomedical community has expended significant resources to discover the specific biological role of members of the PTK family, their function in differentiation processes, their involvement in tumorigenesis and in other diseases, the biochemical mechanisms underlying their signal transduction pathways activated upon ligand stimulation and the development of novel drugs.

Tyrosine kinases can be of the receptor-type (having extracellular, transmembrane and intracellular domains) or the non-receptor type (being wholly intracellular).

**Receptor Tyrosine Kinases.** The RTKs comprise a large family of transmembrane receptors with diverse biological activities. The intrinsic function of RTKs is activated upon ligand binding, which results in phosphorylation of the receptor and multiple cellular substrates, and subsequently in a variety of cellular responses. Ullrich & Schlessinger, 1990, *Cell* 61:203-212.

At present, at least nineteen (19) distinct RTK subfamilies have been identified. One RTK subfamily, designated the HER subfamily, is believed to be comprised of EGFR, HER2, HER3 and HER4. Ligands to the Her subfamily of receptors include epithelial growth factor (EGF), TGF- $\alpha$ , amphiregulin, HB-EGF, betacellulin and heregulin.

A second family of RTKs, designated the insulin subfamily, is comprised of the INS-R, the IGF-1R and the IR-R. A third family, the "PDGF" subfamily includes the PDGF  $\alpha$  and  $\beta$  receptors, CSF1R, c-kit and FLK-II. Another subfamily of RTKs, identified as the FLK family, is believed to be comprised of the Kinase insert Domain-Receptor fetal liver kinase-1 (KDR/FLK-1), the fetal liver kinase 4 (FLK-4) and the fms-like tyrosine kinase 1 (flt-1). Each of these receptors was initially believed to be receptors for hematopoietic growth factors. Two other subfamilies of RTKs have been designated as the FGF receptor family (FGFR1, FGFR2, FGFR3 and FGFR4) and the Met subfamily (c-met and Ron).

Because of the similarities between the PDGF and FLK subfamilies, the two subfamilies are often considered together. The known RTK subfamilies are identified in Plowman et al., 1994, *DN&P* 7(6):334-339, which is incorporated herein by reference.

**The Non-Receptor Tyrosine Kinases.** The non-receptor tyrosine kinases represent a collection of cellular enzymes which lack extracellular and transmembrane sequences. At

present, over twenty-four individual non-receptor tyrosine kinases, comprising eleven (11) subfamilies (Src, Frk, Btk, Csk, Abl, Zap70, Fes/Fps, Fak, Jak, Ack and LIMK) have been identified. At present, the Src subfamily of non-receptor  
5 tyrosine kinases is comprised of the largest number of PTKs and include Src, Yes, Fyn, Lyn, Lck, Blk, Hck, Fgr and Yrk. The Src subfamily of enzymes has been linked to oncogenesis. A more detailed discussion of non-receptor tyrosine kinases is provided in Bolen, 1993, *Oncogene* 8:2025-2031, which is  
10 incorporated herein by reference.

Many of the tyrosine kinases, whether an RTK or non-receptor tyrosine kinase, have been found to be involved in cellular signaling pathways leading to cellular signal assays signalling pathways leading to pathogenic conditions,  
15 including cancer, psoriasis and hyper immune response.

**Development Of Compounds To Modulate The PTKs.** In view of the surmised importance of PTKs to the control, regulation and modulation of cell proliferation the diseases and disorders associated with abnormal cell proliferation, many  
20 attempts have been made to identify receptor and non-receptor tyrosine kinase "inhibitors" using a variety of approaches, including the use of mutant ligands (U.S. Application No. 4,966,849), soluble receptors and antibodies (Application No. WO 94/10202; Kendall & Thomas, 1994, *Proc. Nat'l Acad. Sci*  
25 90:10705-09; Kim, et al., 1993, *Nature* 362:841-844), RNA ligands (Jellinek, et al., *Biochemistry* 33:10450-56); Takano, et al., 1993, *Mol. Bio. Cell* 4:358A; Kinsella, et al., 1992, *Exp. Cell Res.* 199:56-62; Wright, et al., 1992, *J. Cellular Phys.* 152:448-57) and tyrosine kinase inhibitors (WO  
30 94/03427; WO 92/21660; WO 91/15495; WO 94/14808; U.S. Patent No. 5,330,992; Mariani, et al., 1994, *Proc. Am. Assoc. Canc r Res.* 35:2268).

More recently, attempts have been made to identify small molecules which act as tyrosine kinase inhibitors. For  
35 example, bis monocyclic, bicyclic or heterocyclic aryl compounds (PCT WO 92/20642), vinylene-azaindole derivatives (PCT WO 94/14808) and 1-cyclopropyl-4-pyridyl-quinolones


(U.S. Patent No. 5,330,992) have been described generally as tyrosine kinase inhibitors. Styryl compounds (U.S. Patent No. 5,217,999), styryl-substituted pyridyl compounds (U.S. Patent No. 5,302,606), certain quinazoline derivatives (EP 5 Application No. 0 566 266 A1), seleoindoles and selenides (PCT WO 94/03427), tricyclic polyhydroxylic compounds (PCT WO 92/21660) and benzylphosphonic acid compounds (PCT WO 91/15495) have been described as compounds for use as tyrosine kinase inhibitors for use in the treatment of  
10 cancer.

The identification of effective small compounds which specifically inhibit signal transduction by modulating the activity of receptor and non-receptor tyrosine kinases to regulate and modulate abnormal or inappropriate cell  
15 proliferation is therefore desirable and the object of this invention.

### 3. SUMMARY OF THE INVENTION

The present invention relates to organic molecules  
20 capable of modulating, regulating and/or inhibiting tyrosine kinase signal transduction. Such compounds are useful for the treatment of diseases related to unregulated TKS transduction, including cell proliferative diseases such as cancer, atherosclerosis, arthritis and restenosis and  
25 metabolic diseases such as diabetes.

In one illustrative embodiment, the compounds of the present invention have the formula:



and pharmaceutically acceptable salts thereof, wherein

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$R_3$ ,  $R_5$ , and  $R_6$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$n$  is 0-3;


$X$  is Br, Cl, F or I;

$R$  is H, alkyl or aryl; and

$R'$  is H, alkyl or aryl.

In another illustrative embodiment, the compounds of the present invention have the formula:

(II)



30

and pharmaceutically acceptable salts thereof, wherein

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,



$\text{SO}_2\text{NRR}'$ ,  $\text{SO}_2\text{R}$ ,  $\text{SR}$ ,  $\text{NO}_2$ ,  $\text{NRR}'$ ,  $\text{OH}$ ,  $\text{CN}$ ,  $\text{C(O)R}$ ,  $\text{OC(O)R}$ ,  $\text{NHC(O)R}$ ,  $(\text{CH}_2)_n\text{CO}_2\text{R}$ , and  $\text{CONRR}'$ ;

$\text{R}_2$ ,  $\text{R}_3$ ,  $\text{R}_4$ , and  $\text{R}_5$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $\text{S(O)R}$ ,  $\text{SO}_2\text{NRR}'$ ,  $\text{SO}_2\text{R}$ ,  $\text{SR}$ ,  $\text{NO}_2$ ,  $\text{NRR}'$ ,  $\text{OH}$ ,  $\text{CN}$ ,  $\text{C(O)R}$ ,  $\text{OC(O)R}$ ,  $\text{NHC(O)R}$ ,  $(\text{CH}_2)_n\text{CO}_2\text{R}$ , and  $\text{CONRR}'$ ;

$\text{Ra}$  and  $\text{Rb}$  are each independently selected from the group consisting of  $\text{H}$ , alkyl and  $\text{C(O)R}$ , or  $\text{NRaRb}$  taken together may be a heterocyclic ring of from 3 to 8 atoms optionally substituted at one or more positions with hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $\text{S(O)R}$ ,  $\text{SO}_2\text{NRR}'$ ,  $\text{SO}_2\text{R}$ ,  $\text{SR}$ ,  $\text{NO}_2$ ,  $\text{NRR}'$ ,  $\text{OH}$ ,  $\text{CN}$ ,  $\text{C(O)R}$ ,  $\text{OC(O)R}$ ,  $\text{NHC(O)R}$ ,  $(\text{CH}_2)_n\text{CO}_2\text{R}$ , and  $\text{CONRR}'$ ;


$n$  is 0-3;

$\text{X}$  is  $\text{Br}$ ,  $\text{Cl}$ ,  $\text{F}$  or  $\text{I}$ ;

$\text{R}$  is  $\text{H}$ , alkyl or aryl; and

$\text{R}'$  is  $\text{H}$ , alkyl or aryl.

In yet another illustrative embodiment, the compounds of the present invention have the formula:



and pharmaceutically acceptable salts thereof, wherein

$\text{R}_1$  is  $\text{H}$  or alkyl;

$\text{R}_2$  is  $\text{O}$  or  $\text{S}$ ;

$\text{R}_3$  is hydrogen;

$\text{R}_4$ ,  $\text{R}_5$ ,  $\text{R}_6$ , and  $\text{R}_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl,

aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';

A is a five membered heteroaryl ring selected from the group consisting of thiophene, pyrrole, pyrazole, imidazole, 1,2,3-triazole, 1,2,4-triazole, oxazole, isoxazole, thiazole, isothiazole, 2-sulfonylfuran, 4-alkylfuran, 1,2,3-oxadiazole, 1,2,4-oxadiazole, 1,2,5-oxadiazole, 1,3,4-oxadiazole, 1,2,3,4-oxatriazole, 1,2,3,5-oxatriazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,3,4-thiatriazole, 1,2,3,5-thiatriazole, and tetrazole, optionally substituted at one or more positions with alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R or CONRR';

n is 0-3;

X is Br, Cl, F or I;

R is H, alkyl or aryl; and


R' is H, alkyl or aryl.

20

In still another illustrative embodiment, the compounds of the present invention have the formula:

25

(IV)



30

and pharmaceutically acceptable salts thereof, wherein:

R<sub>1</sub> is H or alkyl;

R<sub>2</sub> is O or S;

35 R<sub>3</sub> is hydrogen;

R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>7</sub> are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl,

aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';

R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, and R<sub>6</sub> are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';

n is 0-3;

10 X is Br, Cl, F or I;


R is H, alkyl or aryl; and

R' is H, alkyl or aryl.

In a final illustrative embodiment, the compounds of the 15 present invention have the formula:

(V)

20



25 and pharmaceutically acceptable salts thereof, wherein:

R<sub>1</sub> is H or alkyl;

R<sub>2</sub> is O or S;

R<sub>3</sub> is hydrogen;

R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>7</sub> are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R and CONRR';

R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub>, and R<sub>11</sub> are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R,

$\text{SO}_2\text{NRR}'$ ,  $\text{SO}_2\text{R}$ ,  $\text{SR}$ ,  $\text{NO}_2$ ,  $\text{NRR}'$ ,  $\text{OH}$ ,  $\text{CN}$ ,  $\text{C(O)R}$ ,  $\text{OC(O)R}$ ,  $\text{NHC(O)R}$ ,  
 $(\text{CH}_2)_n\text{CO}_2\text{R}$  and  $\text{CONRR}'$ ;

$n$  is 0-3;

$Z$  is  $\text{Br}$ ,  $\text{Cl}$ ,  $\text{F}$ ,  $\text{I}$ , methyl, ethyl,  $n$ -propyl, iso-propyl,  
5  $n$ -butyl, iso-butyl or tert-butyl;

$R$  is  $\text{H}$ , alkyl or aryl; and

$R'$  is  $\text{H}$ , alkyl or aryl.

The present invention is further directed to  
10 pharmaceutical compositions comprising a pharmaceutically  
effective amount of the above-described compounds of formulae  
I-V and a pharmaceutically acceptable carrier or excipient.  
Such a composition is believed to modulate signal  
transduction by a tyrosine kinase, either by inhibition of  
15 catalytic activity, affinity to ATP or ability to interact  
with a substrate.

More particularly, the compositions of the present  
invention may be included in methods for treating diseases  
comprising proliferation, fibrotic or metabolic disorders,  
20 for example cancer, fibrosis, psoriasis, atherosclerosis,  
arthritis, and other disorders related to abnormal  
vasculogenesis and/or angiogenesis, such as diabetic  
retinopathy.

#### 25 4. DETAILED DESCRIPTION OF THE INVENTION

##### 4.1. Definitions

"Pharmaceutically acceptable salt" refers to those salts  
which retain the biological effectiveness and properties of  
the free bases and which are obtained by reaction with  
30 inorganic acids such as hydrochloric acid, hydrobromic acid,  
sulfuric acid, nitric acid, phosphoric acid, methanesulfonic  
acid, ethanesulfonic acid,  $p$ -toluenesulfonic acid, salicylic  
acid and the like.

"Alkyl" refers to a straight-chain, branched or cyclic  
35 saturated aliphatic hydrocarbon. Preferably, the alkyl group  
has 1 to 12 carbons. More preferably, it is a lower alkyl of  
from 1 to 7 carbons, more preferably 1 to 4 carbons. Typical

alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tertiary butyl, pentyl, hexyl and the like. The alkyl group may be optionally substituted with one or more substituents are selected from the group consisting of 5 hydroxyl, cyano, alkoxy, =O, =S, NO<sub>2</sub>, halogen, N(CH<sub>3</sub>), amino, and SH.

"Alkenyl" refers to a straight-chain, branched or cyclic unsaturated hydrocarbon group containing at least one carbon-carbon double bond. Preferably, the alkenyl group has 1 to 10 12 carbons. More preferably it is a lower alkenyl of from 1 to 7 carbons, more preferably 1 to 4 carbons. The alkenyl group may be optionally substituted with one or more substituents selected from the group consisting of hydroxyl, cyano, alkoxy, =O, =S, NO<sub>2</sub>, halogen, N(CH<sub>3</sub>), amino, and SH.

15 "Alkynyl" refers to a straight-chain, branched or cyclic unsaturated hydrocarbon containing at least one carbon-carbon triple bond. Preferably, the alkynyl group has 1 to 12 carbons. More preferably it is a lower alkynyl of from 1 to 7 carbons, more preferably 1 to 4 carbons. The alkynyl group 20 may be optionally substituted with one or more substituents selected from the group consisting of hydroxyl, cyano, alkoxy, =O, =S, NO<sub>2</sub>, halogen, N(CH<sub>3</sub>), amino, and SH.

"Alkoxy" refers to an "-Oalkyl" group.

"Aryl" refers to an aromatic group which has at least 25 one ring having a conjugated pi electron system and includes carbocyclic aryl, heterocyclic aryl and biaryl groups. The aryl group may be optionally substituted with one or more substituents selected from the group consisting of halogen, trihalomethyl, hydroxyl, SH, OH, NO<sub>2</sub>, amine, thioether, cyano, 30 alkoxy, alkyl, and amino.

"Alkaryl" refers to an alkyl that is covalently joined to an aryl group. Preferably, the alkyl is a lower alkyl.

"Carbocyclic aryl" refers to an aryl group wherein the ring atoms are carbon.

35 "Heterocyclic aryl" refers to an aryl group having from 1 to 3 heteroatoms as ring atoms, the remainder of the ring atoms being carbon. Heteroatoms include oxygen, sulfur, and

nitrogen. Thus, heterocyclic aryl groups include furanyl, thienyl, pyridyl, pyrrolyl, N-lower alkyl pyrrolo, pyrimidyl, pyrazinyl, imidazolyl and the like.

"Amide" refers to  $-C(O)-NH-R$ , where R is alkyl, aryl, 5 alkylaryl or hydrogen.

"Thioamide" refers to  $-C(S)-NH-R$ , where R is alkyl, aryl, alkylaryl or hydrogen.

"Amine" refers to a  $-N(R')R''$  group, where R' and R'' are independently selected from the group consisting of 10 alkyl, aryl, and alkylaryl.

"Thioether" refers to  $-S-R$ , where R is alkyl, aryl, or alkylaryl.

"Sulfonyl" refers to  $-S(O)_2-R$ , where R is aryl,  $C(CN)=C$ -aryl,  $CH_2CN$ , alkyaryl, sulfonamide, NH-alkyl, NH-alkylaryl, or 15 NH-aryl.

#### 4.2. The Invention

The present invention relates to compounds capable of regulating and/or modulating tyrosine kinase signal 20 transduction and more particularly receptor and non-receptor tyrosine kinase signal transduction.

Receptor tyrosine kinase mediated signal transduction is initiated by extracellular interaction with a specific growth factor (ligand), followed by receptor dimerization, transient 25 stimulation of the intrinsic protein tyrosine kinase activity and phosphorylation. Binding sites are thereby created for intracellular signal transduction molecules and lead to the formation of complexes with a spectrum of cytoplasmic signalling molecules that facilitate the appropriate cellular 30 response (e.g., cell division, metabolic effects to the extracellular microenvironment). See, Schlessinger and Ullrich, 1992, *Neuron* 9:303-391.

It has been shown that tyrosine phosphorylation sites in growth factor receptors function as high-affinity binding 35 sites for SH2 (src homology) domains of signaling molecules. Fantl et al., 1992, *Cell* 69:413-423; Songyang et al., 1994, *Mol. Cell. Biol.* 14:2777-2785; Songyang et al., 1993, *Cell*

72:767-778; and Koch et al., 1991, *Science* 252:668-678. Several intracellular substrate proteins that associate with receptor tyrosine kinases have been identified. They may be divided into two principal groups: (1) substrates which have  
5 a catalytic domain; and (2) substrates which lack such domain but serve as adapters and associate with catalytically active molecules. Songyang et al., 1993, *Cell* 72:767-778. The specificity of the interactions between receptors and SH2 domains of their substrates is determined by the amino acid  
10 residues immediately surrounding the phosphorylated tyrosine residue. Differences in the binding affinities between SH2 domains and the amino acid sequences surrounding the phosphotyrosine residues on particular receptors are consistent with the observed differences in their substrate  
15 phosphorylation profiles. Songyang et al., 1993, *Cell* 72:767-778. These observations suggest that the function of each receptor tyrosine kinase is determined not only by its pattern of expression and ligand availability but also by the array of downstream signal transduction pathways that are  
20 activated by a particular receptor. Thus, phosphorylation provides an important regulatory step which determines the selectivity of signaling pathways recruited by specific growth factor receptors, as well as differentiation factor receptors.

25 Tyrosine kinase signal transduction results in, among other responses, cell proliferation, differentiation and metabolism. Abnormal cell proliferation may result in a wide array of disorders and diseases, including the development of neoplasia such as carcinoma, sarcoma, leukemia, glioblastoma,  
30 hemangioma, psoriasis, arteriosclerosis, arthritis and diabetic retinopathy (or other disorders related to uncontrolled angiogenesis and/or vasculogenesis).

This invention is therefore directed to compounds which regulate, modulate and/or inhibit tyrosine kinase signal  
35 transduction by affecting the enzymatic activity of the RTKs and/or the non-receptor tyrosine kinases and interfering with the signal transduced such proteins. More particularly, the

present invention is directed to compounds which regulate, modulate and/or inhibit the RTK and/or non-receptor tyrosine kinase mediated signal transduction pathways as a therapeutic approach to cure many kinds of solid tumors, including but not limited to carcinoma, sarcoma, leukemia, erythroblastoma, glioblastoma, meningioma, astrocytoma, melanoma and myoblastoma. Indications may include, but are not limited to brain cancers, bladder cancers, ovarian cancers, gastric cancers, pancreas cancers, colon cancers, blood cancers, lung cancers and bone cancers.


#### 4.3. The Compounds

In one embodiment, the invention provides compounds having the formula:

15

(I)

20



and the pharmaceutically acceptable salts thereof, wherein

25  $R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, 30 aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_3R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$R_3$ ,  $R_5$ , and  $R_6$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, 35 alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_3R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;



n is 0-3;  
X is Br, Cl, F or I; and  
R is H, alkyl or aryl; and  
R' is H, alkyl or aryl.

5

In a preferred embodiment of the compounds of formula I, R<sub>3</sub>' is selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where R is H, alkyl or aryl; and R<sub>4</sub>' is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where R is H, alkyl or aryl.

A particularly preferred compound of formula I is 3-(2-chloro-4-hydroxybenzylidenenyl)-2-indolinone (SU4932).


15

In another embodiment, the invention provides compounds having the formula:

20

(II)

25



and the pharmaceutically acceptable salts thereof, wherein

R<sub>1</sub> is H or alkyl;

R<sub>2</sub> is O or S;

30 R<sub>3</sub> is hydrogen;

R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>7</sub> are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, 35 (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';

R<sub>2</sub>', R<sub>3</sub>', R<sub>4</sub>', and R<sub>5</sub>' are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl,

aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';

Ra and Rb are each independently selected from the group  
 5 consisting of H, alkyl and C(O)R, or NRaRb taken together may be a heterocyclic ring of from 3 to 8 atoms optionally substituted at one or more positions with hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN,  
 10 C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, or CONRR';  
 n is 0-3;  
 X is Br, Cl, F or I; and  
 R is H, alkyl or aryl; and  
 R' is H, alkyl or aryl.

15

In a preferred embodiment of the compounds of formula II, R<sub>3</sub>' is selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where R is H, alkyl or aryl; and R<sub>3</sub>' is  
 20 selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where R is H, alkyl or aryl.


A particularly preferred compound of formula II is 3-(4-Dimethylaminobenzylidenyl)-2-indolinone (SU4312).

25

In yet another embodiment, the invention provides compounds having the formula:

30

(III)



35 and the pharmaceutically acceptable salts thereof, wherein  
 R<sub>1</sub> is H or alkyl;  
 R<sub>2</sub> is O or S;

R<sub>3</sub> is hydrogen;

R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>7</sub> are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';

A is a five membered heteroaryl ring selected from the group consisting of thiophene, pyrrole, pyrazole, imidazole, 1,2,3-triazole, 1,2,4-triazole, oxazole, isoxazole, thiazole, isothiazole, 2-sulfonylfuran, 4-alkylfuran, 1,2,3-oxadiazole, 1,2,4-oxadiazole, 1,2,5-oxadiazole, 1,3,4-oxadiazole, 1,2,3,4-oxatriazole, 1,2,3,5-oxatriazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,3,4-thiatriazole, 1,2,3,5-thiatriazole, and tetrazole, optionally substituted at one or more positions with alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R or CONRR';

n is 0-3;

X is Br, Cl, F or I;

R is H, alkyl or aryl; and

R' is H, alkyl or aryl.

In a preferred embodiment of the invention, the compound of formula III is 3-[(2,3-Dimethylpyrrol-5-yl)methylene]-2-indolinone (SU5416).

In still another embodiment, the invention provides compounds having the formula:

(IV)


and the  
pharmaceutically  
acceptable salts  
thereof, wherein:

5         $R_1$  is H or  
alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and



10  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

15         $R_1$ ,  $R_4$ ,  $R_5$ , and  $R_6$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

20         $n$  is 0-3;

$X$  is Br, Cl, F or I;

$R$  is H, alkyl or aryl; and


$R'$  is H, alkyl or aryl.

25        In a preferred embodiment of the compound of formula IV,  $R_3$  is selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where  $R$  is H, alkyl or aryl; and  $R_5$  is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where  $R$  is H, alkyl or aryl.

A particularly preferred compound of formula IV is 3-(2-Ethoxybenzylidenyl)-2-indolinone (SU5204).

In a final embodiment, the invention provides compounds  
35 having the formula:

(V)



5

and the  
pharmaceutically  
acceptable salts  
thereof, wherein:

10  $R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from  
the group consisting of hydrogen, alkyl, alkoxy, aryl,  
15 aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  
 $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  
 $(CH_2)_nCO_2R$  and  $CONRR'$ ;

$R_2'$ ,  $R_3'$ ,  $R_5'$ , and  $R_6'$  are each independently selected from  
the group consisting of hydrogen, alkyl, alkoxy, aryl,  
20 aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  
 $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  
 $(CH_2)_nCO_2R$  and  $CONRR'$ ;

$n$  is 0-3;

$Z$  is Br, Cl, F, I, methyl, ethyl, n-propyl, iso-propyl,  
25 n-butyl, iso-butyl or tert-butyl;

$R$  is H, alkyl or aryl; and

$R'$  is H, alkyl or aryl.

In a preferred embodiment of the compounds of formula V,  
 $R_3'$  is selected from the group consisting of methyl, ethyl, n-  
30 propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen,  
aryl and OR, where  $R$  is H, alkyl or aryl; and  $R_5'$  is selected  
from the group consisting of hydrogen, methyl, ethyl, n-  
propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen,  
aryl and OR, where  $R$  is H, alkyl or aryl.

35 A particularly preferred compound of formula V is 3-(4-  
Bromobenzylidenyl)-2-indolinone (SU4942).

The chemical formulae referred herein may exhibit the phenomena of tautomerism or structural isomerism. For example, the compounds described herein may be adopt a *cis* or *trans* conformation about the double bond connecting the indolinone 3-substituent to the indolinone ring, or may be mixtures of *cis* and *trans* isomers. As the formulae drawing within this specification can only represent one possible tautomeric or structural isomeric form, it should be understood that the invention encompasses any tautomeric or structural isomeric form, or mixtures thereof, which possesses the ability to regulate, inhibit and/or modulate tyrosine kinase signal transduction or cell proliferation and is not limited to any one tautomeric or structural isomeric form utilized within the formulae drawing.

In addition to the above-described compounds and their pharmaceutically acceptable salts, the invention is further directed, where applicable, to solvated as well as unsolvated forms of the compounds (e.g. hydrated forms) having the ability to regulate and/or modulate cell proliferation.

The compounds described herein may be prepared by any process known to be applicable to the preparation of chemically-related compounds. Suitable processes are illustrated in the examples. Necessary starting materials may be obtained by standard procedures of organic chemistry.

An individual compound's relevant activity and efficacy as an agent to affect receptor tyrosine kinase mediated signal transduction may be determined using available techniques. Preferentially, a compound is subjected to a series of screens to determine the compound's ability to modulate, regulate and/or inhibit cell proliferation. These screens, in the order in which they are conducted, include biochemical assays, cell growth assays and *in vivo* experiments.

35

#### 4.4. Indications

The compounds described herein are useful for treating disorders related to unregulated tyrosine kinase signal transduction, including cell proliferative disorders, fibrotic disorders and metabolic disorders.

Cell proliferative disorders which can be treated or further studied by the present invention include cancers, blood vessel proliferative disorders and mesangial cell proliferative disorders.

10 Blood vessel proliferative disorders refer to angiogenic and vasculogenic disorders generally resulting in abnormal proliferation of blood vessels. The formation and spreading of blood vessels, or vasculogenesis and angiogenesis, respectively, play important roles in a variety of physiological processes such as embryonic development, corpus luteum formation, wound healing and organ regeneration. They also play a pivotal role in cancer development. Other examples of blood vessel proliferation disorders include arthritis, where new capillary blood vessels invade the joint and destroy cartilage, and ocular diseases, like diabetic retinopathy, where new capillaries in the retina invade the vitreous, bleed and cause blindness. Conversely, disorders related to the shrinkage, contraction or closing of blood vessels, such as restenosis, are also implicated.

25 Fibrotic disorders refer to the abnormal formation of extracellular matrix. Examples of fibrotic disorders include hepatic cirrhosis and mesangial cell proliferative disorders. Hepatic cirrhosis is characterized by the increase in extracellular matrix constituents resulting in the formation of a hepatic scar. Hepatic cirrhosis can cause diseases such as cirrhosis of the liver. An increased extracellular matrix resulting in a hepatic scar can also be caused by viral infection such as hepatitis. Lipocytes appear to play a major role in hepatic cirrhosis. Other fibrotic disorders implicated include atherosclerosis (see, below).

35 Mesangial cell proliferative disorders refer to disorders brought about by abnormal proliferation of

mesangial cells. Mesangial proliferative disorders include various human renal diseases, such as glomerulonephritis, diabetic nephropathy, malignant nephrosclerosis, thrombotic microangiopathy syndromes, transplant rejection, and  
5 glomerulopathies. The PDGF-R has been implicated in the maintenance of mesangial cell proliferation. Floege et al., 1993, *Kidney International* 43:47S-54S.

PTKs have been associated with such cell proliferative disorders. For example, some members of the RTK family have  
10 been associated with the development of cancer. Some of these receptors, like the EGFR (Tuzi et al., 1991, *Br. J. Cancer* 63:227-233; Torp et al., 1992, *APMIS* 100:713-719) HER2/neu (Slamon et al., 1989, *Science* 244:707-712) and the PDGF-R (Kumabe et al., 1992, *Oncogene* 7:627-633) are  
15 overexpressed in many tumors and/or persistently activated by autocrine loops. In fact, in the most common and severe cancers these receptor overexpressions (Akbasak and Suner-Akbasak et al., 1992, *J. Neurol. Sci.* 111:119-133; Dickson et al., 1992, *Cancer Treatment Res.* 61:249-273; Korc et al.,  
20 1992, *J. Clin. Invest.* 90:1352-1360) and autocrine loops (Lee and Donoghue, 1992, *J. Cell. Biol.* 118:1057-1070; Korc et al., *supra*; Akbasak and Suner-Akbasak et al., *supra*) have been demonstrated. For example, the EGFR receptor has been associated with squamous cell carcinoma, astrocytoma,  
25 glioblastoma, head and neck cancer, lung cancer and bladder cancer. HER2 has been associated with breast, ovarian, gastric, lung, pancreas and bladder cancer. The PDGF-R has been associated with glioblastoma, lung, ovarian, melanoma and prostate. The RTK c-met has been generally associated  
30 with hepatocarcinogenesis and thus hepatocellular carcinoma. Additionally, c-met has been linked to malignant tumor formation. More specifically, the RTK c-met has been associated with, among other cancers, colorectal, thyroid, pancreatic and gastric carcinoma, leukemia and lymphoma.  
35 Additionally, over-expression of the c-met gene has been detected in patients with Hodgkins disease, Burkitts disease, and the lymphoma cell line.



The IGF-IR, in addition to being implicated in nutritional support and in type-II diabetes, has also been associated with several types of cancers. For example, IGF-I has been implicated as an autocrine growth stimulator for several tumor types, e.g. human breast cancer carcinoma cells (Arteaga et al., 1989, *J. Clin. Invest.* 84:1418-1423) and small lung tumor cells (Macauley et al., 1990, *Cancer Res.* 50:2511-2517). In addition, IGF-I, integrally involved in the normal growth and differentiation of the nervous system, appears to be an autocrine stimulator of human gliomas. Sandberg-Nordqvist et al., 1993, *Cancer Res.* 53:2475-2478. The importance of the IGF-IR and its ligands in cell proliferation is further supported by the fact that many cell types in culture (fibroblasts, epithelial cells, smooth muscle cells, T-lymphocytes, myeloid cells, chondrocytes, osteoblasts, the stem cells of the bone marrow) are stimulated to grow by IGF-I. Goldring and Goldring, 1991, *Eukaryotic Gene Expression* 1:301-326. In a series of recent publications, Baserga even suggests that IGF-I-R plays a central role in the mechanisms of transformation and, as such, could be a preferred target for therapeutic interventions for a broad spectrum of human malignancies. Baserga, 1995, *Cancer Res.* 55:249-252; Baserga, 1994, *Cell* 79:927-930; Coppola et al., 1994, *Mol. Cell. Biol.* 14:4588-4595.

The association between abnormalities in RTKs and disease are not restricted to cancer, however. For example, RTKs have been associated with metabolic diseases like psoriasis, diabetes mellitus, wound healing, inflammation, and neurodegenerative diseases. For example, the EGF-R is indicated in corneal and dermal wound healing. Defects in the Insulin-R and the IGF-1R are indicated in type-II diabetes mellitus. A more complete correlation between specific RTKs and their therapeutic indications is set forth in Plowman et al., 1994, *DN&P* 7:334-339.

Not only receptor type tyrosine kinases, but also many cellular tyrosine kinases (CTKs) including src, abl, fps,

yes, fyn, lyn, lck, blk, hck, fgr, yrk (reviewed by Bolen et al., 1992, *FASEB J.* 6:3403-3409) are involved in the proliferative and metabolic signal transduction pathway and thus in indications of the present invention. For example, 5 mutated src (v-src) has been demonstrated as an oncoprotein (pp60<sup>v-src</sup>) in chicken. Moreover, its cellular homolog, the proto-oncogene pp60<sup>c-src</sup> transmits oncogenic signals of many receptors. For example, overexpression of EGF-R or HER2/neu in tumors leads to the constitutive activation of pp60<sup>c-src</sup>, 10 which is characteristic for the malignant cell but absent from the normal cell. On the other hand, mice deficient for the expression of c-src exhibit an osteopetrotic phenotype, indicating a key participation of c-src in osteoclast function and a possible involvement in related disorders. 15 Similarly, Zap 70 is implicated in T-cell signalling.

Furthermore, the identification of CTK modulating compounds to augment or even synergize with RTK aimed blockers is an aspect of the present invention.

Finally, both RTKs and non-receptor type kinases have 20 been connected to hyperimmune disorders.

#### 4.5. Pharmaceutical Formulations And Routes Of Administration

The compounds described herein can be administered to a human patient *per se*, or in pharmaceutical compositions where 25 it is mixed with suitable carriers or excipient(s). Techniques for formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition.

##### 30 4.5.1. Routes Of Administration.

Suitable routes of administration may, for example, include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as 35 intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound directly into a solid tumor, often in a depot or sustained release formulation.

5 Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with tumor-specific antibody. The liposomes will be targeted to and taken up selectively by the tumor.

#### 4.5.2. Composition/Formulation.

10 The pharmaceutical compositions of the present invention may be manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

15 Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which  
20 can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution,  
25 Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated  
30 readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral  
35 ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained solid excipient, optionally grinding a resulting mixture, and processing the

mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered

in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, .g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable 5 gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as 10 lactose or starch.

The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi- 15 dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

20 Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or 25 vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. 30 Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Alternatively, the active ingredient may be in powder 35 form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

- 5 In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the
- 10 compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.
- 15 A pharmaceutical carrier for the hydrophobic compounds of the invention is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The cosolvent system may be the VPD co-solvent system. VPD is a solution of 3% w/v
- 20 benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:5W) consists of VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic
- 25 compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied: for
- 30 example, other low-toxicity nonpolar surfactants may be used instead of polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, e.g. polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose.
- 35 Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or

carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, 5 such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various of sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for 10 a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

The pharmaceutical compositions also may comprise 15 suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

20 Many of the PTK modulating compounds of the invention may be provided as salts with pharmaceutically compatible counterions. Pharmaceutically compatible salts may be formed with many acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. 25 Salts tend to be more soluble in aqueous or other protonic solvents that are the corresponding free base forms.

#### 4.5.3. Effective Dosage.

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active 30 ingredients are contained in an amount effective to achieve its intended purpose. More specifically, a therapeutically effective amount means an amount of compound effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated. 35 Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

For any compound used in the methods of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes the  $IC_{50}$  as determined in cell culture (i.e., the concentration of the test compound which achieves a half-maximal inhibition of the PTK activity). Such information can be used to more accurately determine useful doses in humans.

10 Toxicity and therapeutic efficacy of the compounds described herein can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the  $LD_{50}$  (the dose lethal to 50% of the population) and the  $ED_{50}$  (the dose therapeutically effective  
15 in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between  $LD_{50}$  and  $ED_{50}$ . Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies  
20 can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the  $ED_{50}$  with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of  
25 administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See e.g., Fingl et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1).

30 Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the kinase modulating effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data; e.g.,  
35 the concentration necessary to achieve 50-90% inhibition of the kinase using the assays described herein. Dosages necessary to achieve the MEC will depend on individual



characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations.

Dosage intervals can also be determined using MEC value. 5 Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%.

In cases of local administration or selective uptake, the effective local concentration of the drug may not be 10 related to plasma concentration.

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

#### 15 4.5.4. Packaging

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister 20 pack. The pack or dispenser device may be accompanied by instructions for administration. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labelled for treatment of an 25 indicated condition. Suitable conditions indicated on the label may include treatment of a tumor, inhibition of angiogenesis, treatment of fibrosis, diabetes, and the like.

#### 5. EXAMPLE: Compound Synthesis

30 The compounds of the present invention may be synthesized according to known techniques. The following represent preferred methods for synthesizing the compounds of the claimed invention.

##### 35 5.1. General Syntheses of 3-Substituted-2-Indolinone Analogs (SU4312 and SU4314 Analogs)

The following general methodologies were used to synthesize 3-substituted-2-indolinone compounds of the invention.

5.1.1. Method A

5 A reaction mixture of the proper oxindole (2-indolinone) (1 equiv.), the appropriate aldehyde (1.2 equiv.), and piperidine (0.1 equiv.) in ethanol (1 - 2 mL / 1 mmol oxindole) was stirred at 90°C for 3 - 5 h. After cooling, the precipitate was filtered, washed with cold  
10 ethanol, and dried to yield the target compound.

5.1.2. Method B

Preparation of The Proper Aldehydes via Vilsmeier Reaction. To a solution of N,N-dimethylformamide (1.2  
15 equiv.) in 1,2-dichloroethane (2.0 mL / 1.0 mmole of starting material) was added dropwise phosphorus oxychloride (1.2 equiv.) at 0°C. The ice-bath was removed and the reaction mixture was further stirred for 30 min. The proper starting material (1.0 equiv.) was added to the above solution  
20 portionwise and the reaction mixture was stirred at 50-70°C for 5 h - 2 days. The reaction mixture was poured into ice-cold 1N sodium hydroxide solution (pH = 9 after mixing) and the resulting mixture was stirred at room temperature for 1 h. The organic layer was separated and the aqueous layer was  
25 extracted with ethyl acetate. The combined organic layer was washed with brine until pH = 7, dried over anhydrous sodium sulfate and evaporated. The residue was chromatographed on a silica gel column eluting with a solvent mixture of ethyl acetate and hexane to afford the title compound.

30 Synthesis for 3-Substituted-2-Indolinone Analogs. A reaction mixture of the proper oxindole (2-indolinone) (1 equiv.), the appropriate aldehyde (1.2 equiv.), and piperidine (0.1 equiv.) in ethanol (1 - 2 mL / 1 mmol oxindole) was stirred at 90°C for 3 - 5 h. After cooling,  
35 the precipitate was filtered, washed with cold ethanol and dried to yield the target compound.

### 5.2. Synthesis Of 3-Benzylidene-2-Indolinone (SU4928)

The preferred method for synthesizing 3-benzylidene-2-indolinone is as follows: Added 123.2  $\mu$ l of benzaldehyde and 40  $\mu$ l of piperidine to a solution of 137.0 mg of oxindole in 2.0 ml methanol. Reflux the reaction mixture for 3 hours and cool down the mixture in an ice-water bath. Filter the resulting precipitate, wash with cold methanol and dry in an oven at 40°C overnight. Approximately 129.0 mg of the compound was obtained using such protocol.

10

### 5.3. Synthesis Of 3-[(Pyrid-4-yl)methylene]-2-indolinone (SU5212)

The preferred method for synthesizing 3-[(Pyrid-4-yl)methylene]-2-indolinone as follows: Add 117.0  $\mu$ l of 4-pyridinecarboxaldehyde and 40  $\mu$ l of piperidine to a solution of 138.0 mg of oxindole in 2.0 ml methanol. The reaction mixture was refluxed for 3 hours and cooled down in an ice-water bath. The resulting precipitate was filtered, washed with cold methanol and dried in an oven at 40°C overnight to give 134.5 mg of the compound.

20

### 5.4. Synthesis of 3-[4-(morpholin-4-yl)benzylidenyl]-2-indolinone (SU4981) (Method B):

4-(Morpholin-4-yl)benzaldehyde. To a solution of 15 mL of N,N-dimethylformamide in 50 mL of 1,2-dichloroethane was added dropwise 10 mL of phosphorus oxychloride at 0°C. The ice-bath was removed and the reaction mixture was further stirred for 30 min. 4-Phenylmorpholine (16.3 g) was added to the above solution portionwise and the reaction mixture was refluxed for 2 days. Triethylamine (2.5 mL) was added to the above reaction mixture and the reaction was refluxed for 2 day. The reaction mixture was poured into ice-cold 1N sodium hydroxide solution (pH = 9 after mixing) and the resulting mixture was stirred at room temperature for 1 h. The organic layer was separated and the aqueous layer was extracted with 2 x 20 mL of dichloromethane. The combined organic layer was

25

30

35

washed with brine until pH = 7, dried over anhydrous sodium sulfate and evaporated. The residue was separated on a silica gel column eluting with a solvent mixture of ethyl acetate and hexane to afford 12.95 g (68%) of the title compound as a white solid.

3-[4-(Morpholin-4-yl)benzylidenyl]-2-indolinone (SU4981). A reaction mixture of 6.66 g of oxindole, 11.50 g of the 4-(morpholine-4-yl)benzaldehyde, and 5 mL of piperidine in 50 mL of ethanol was stirred at 90°C for 5 h. After cooling, the precipitate was filtered, washed with cold ethanol, and dried to yield 15.0 g (98%) of the title compound as a yellow solid.

5.5. Synthesis of 3-[4-(4-Formylpiperazin-1-yl)benzylidenyl]-2-indolinone (SU4984) (Method B):  
4-(4-Formylpiperazin-1-yl)benzaldehyde. To a solution of 3.9 mL (30 mmoles) of N,N-dimethylformamide in 20 mL of 1,2-dichloroethane was added dropwise 3.0 mL (3.9 mmoles) of phosphorus oxychloride at 0°C. The ice-bath was removed and the reaction mixture was further stirred for 15 min. 1-Phenylpiperazine (16.0 g, 10 mmoles) was added to the above solution portionwise and the reaction mixture was stirred at 50°C for 1 h. The reaction mixture was poured into ice-cold 1N sodium hydroxide solution and stirred at room temperature for 1 h. The organic layer was separated and the aqueous layer was extracted with 2 x 20 mL of ethyl acetate. The combined organic layer was washed with brine until pH = 7, dried over anhydrous sodium sulfate and evaporated. The residue was separated on a silica gel column eluting with a mixture of ethyl acetate and hexane to afford 9.0 g (41%) of the title compound a light yellow solid.

3-[4-(4-Formylpiperazin-1-yl)benzylidenyl]-2-indolinone (SU4984). A reaction mixture of 133.15 mg of oxindole, 228.3 mg of 4-(piperazin-1-yl)benzaldehyde, and 3 drops of piperidine in 2 mL of ethanol was stirred at 90°C for 5 h. After cooling, the precipitate was filtered, washed with cold

ethanol and dried to yield 199.5 mg (65%) of the title compound a yellow solid.

5.6. Synthesis of 3-[4-(Piperidin-1-yl)benzylidenyl]-2-indolinone (SU5450) (Method B).

5 4-(Piperidin-1-yl)benzaldehyde. To a solution of 2.3 mL (30 mmoles) of N,N-dimethylformamide in 10 mL of 1,2-dichloroethane was added dropwise 2.8 mL (30 mmoles) of phosphorus oxychloride at 0°C. The ice-bath was removed and the reaction mixture was stirred for 15 min. 1-Phenylpiperidine (3.2 mL, 20 mmoles) was added to the above solution portionwise and the reaction mixture was refluxed overnight. The reaction mixture was poured into ice-cold 2N sodium hydroxide solution and stirred at room temperature for 1 h. The organic layer was separated and the aqueous layer was extracted with 2 x 20 mL of ethyl acetate. The combined organic layer was washed with brine until pH = 7, dried over anhydrous sodium sulfate and evaporated. The residue was separated on a silica gel column eluting with ethyl acetate and hexane to afford 1.5 g (40%) of the title compound as a white solid.

3-[4-(Piperidin-1-yl)benzylidenyl]-2-indolinone (SU5450). A reaction mixture of 134.0 mg of oxindole, 226.8 g of 4-(piperidine-1-yl)benzaldehyde, and 3 drops of piperidine in 2 mL of ethanol was stirred at 90°C for 5 h. After cooling, the precipitate was filtered, washed with cold ethanol, and dried to yield 268.5 mg (88%) of the title compound as a yellow solid.

5.7. Synthesis of 3-[2-Chloro-4-methoxybenzylidenyl]-2-indolinone (SU5480).

2-Chloro-4-methoxybenzaldehyde. The reaction mixture of 1.0 g (6.4 mmoles) of 2-chloro-4-hydroxybenzaldehyde, 4.4 g (32 mmoles) of potassium carbonate, and 1.4 g (9.6 mmoles) of methyl iodide in 10 mL of N,N-dimethylformamide was stirred at 70°C for 2 h and poured into ice water. The precipitate was filtered, washed with water, and dried at 40°C in vacuum

oven overnight to yield 750 mg (68%) of the title compound as a light pink solid.

3-[2-Chloro-4-methoxybenzylidenyl]-2-indolinone (SU5480). The reaction mixture of 487.9 mg (3.7 mmoles) of oxindole, 750 mg (4.3 mmoles) of 2-chloro-4-methoxybenzaldehyde and 4 drops of piperidine in 5 mL of ethanol was heated to 90°C for 2 h and cooled to room temperature. The yellow precipitate was filtered, washed with cold ethanol, and dried at 40°C in a vacuum oven overnight to give 680.2 mg (62%) of the title compound.

5.8. Synthesis of 3-[(4-Methylthien-2-yl)methylene]-2-indolinone (SU5401).

A reaction mixture of 133.0 mg of oxindole, 151.2 mg of the 4-methylthiophene-2-carboxaldehyde, and 3 drops of piperidine in 3 mL of ethanol was stirred at 90°C for 3 h. After cooling, the precipitate was filtered, washed with cold ethanol, and dried to yield 147.3 mg (61%) of the title compound as a yellow solid.

20

5.9. Synthesis of 3-[(3-Methylpyrrol-2-yl)methylene]-2-indolinone (SU5404).

A reaction mixture of 133.0 mg of oxindole, 130.9 mg of the 3-methylpyrrole-2-carboxaldehyde, and 3 drops of piperidine in 2 mL of ethanol was stirred at 90°C for 3 h. After cooling, the precipitate was filtered, washed with cold ethanol, and dried to yield 150.9 mg (67%) of the title compound as a yellow solid.

5.10. Synthesis of 3-[(3,4-Dimethylpyrrol-2-yl)methylene]-2-indolinone (SU5406)

3-[(3,4-Dimethylpyrrol-2-yl)methylene]-2-indolinone was synthesized as described in J. Heterocyclic Chem. 13:1145-1147 (1976).

Ethyl 4-methylpyrrol-3-carboxylate. A solution of 11.86 g (0.1 moles) of ethyl crotonate and 19.50 g (0.1 moles) of p-toluenesulfonylmethylisocyanide in 500 mL of a 2:1 ether/dimethylsulfoxide was added dropwise into a suspension

of 6.8 g of sodium hydride (60% mineral oil dispersion, 0.17 moles) in ether at room temperature. Upon completion of addition the reaction mixture was stirred for 30 min and dilute with 400 mL of water. The aqueous layer was extracted 5 with 3x100 mL of ether. The combined ether extracts were passed through a column of alumina eluting with dichloromethane. The organic solvent was evaporated and the resulting residue was solidified on standing. The solid was washed with hexane and dried at 40°C in vacuum oven overnight 10 to yield 12.38 g (80%) of the title compound.

Preparation of 3,4-Dimethylpyrrole. To a solution of 23 g (80 mmoles) of sodium dihydrobis(2-methoxyethoxy aluminate) was added dropwise of a solution of 5 g (34 mmoles) of ethyl 4-methylpyrrol-3-carboxylate in 50 mL of benzene at room 15 temperature under nitrogen atmosphere. The reaction mixture was stirred for 18 h. Water (100 mL) was added to the reaction mixture. The organic layer was separated, washed with brine and dried over anhydrous sodium sulfate. The solvent was removed and the residue was distilled giving 1.2 20 g (44%) of the title compound.

Preparation of 3,4-Dimethylpyrrole-2-carboxaldehyde. T a solution of 0.92 mL (12 mmoles) of N,N-dimethylformamide in 10 mL of 1,2-dichloroethane was added dropwise 1.0 mL (12 mmoles) of phosphorus oxychloride at 0°C. The ice-bath was 25 removed and the reaction mixture was further stirred for 30 min. 3,4-Dimethylpyrrole (960.0 mg, 10 mmoles) was added to the above solution portionwise and the reaction mixture was stirred at 50°C for 5 h. The reaction mixture was poured into ice-cold 1N sodium hydroxide solution (pH = 9 after 30 mixing) and the resulting mixture was stirred at room temperature for 1 h. The organic layer was separated and the aqueous layer was extracted with ethyl acetate. The combined organic layer was washed with brine until pH = 7, dried over anhydrous sodium sulfate and evaporated. The residue was 35 chromatographed on a silica gel column eluting with a solvent mixture of ethyl acetate and hexane to afford 610 mg (50%) of the title compound.

3-[(3,4-Dimethylpyrrol-2-yl)methylene]-2-indolinone (SU5406). A reaction mixture of 67.0 mg (0.5 mmoles) of oxindole, 73.0 mg (0.6 mmoles) of the 3,4-dimethylpyrrole-2-carboxaldehyde, and 2 drops of piperidine in 2 mL of ethanol was stirred at 90°C for 3 h. After cooling, the precipitate was filtered, washed with cold ethanol, and dried to yield 87.7 mg (37%) of the title compound as a yellow solid.

5.11. Synthesis of 3-[(2,4-Dimethyl-3-ethoxycarbonylpyrrol-5-yl)methylene]-2-indolinone (SU5408)

A reaction mixture of 134.0 mg of oxindole, 234.3 mg of the 4-ethoxycarbonyl-3,5-dimethylpyrrole-2-carboxaldehyde, and 3 drops of piperidine in 3 mL of ethanol was stirred at 90°C for 3 h. After cooling, the precipitate was filtered, washed with cold ethanol, and dried to yield 244.6 mg (79%) of the title compound as a yellow solid.

5.12. Synthesis of 3-[(2,3-Dimethylpyrrol-5-yl)methylene]-2-indolinone (SU5416)

A reaction mixture of 134.0 mg of oxindole, 147.8 mg of the 3,5-dimethylpyrrole-2-carboxaldehyde, and 3 drops of piperidine in 2 mL of ethanol was stirred at 90°C for 3 h. After cooling, the precipitate was filtered, washed with cold ethanol, and dried to yield 136.7 mg (57%) of the title compound as a yellow solid.

5.13. Synthesis of 3-[(2-Methylmercaptothiophen-5-yl)methylene]-2-indolinone (SU5419)

A reaction mixture of 134.0 mg of oxindole, 189.9 mg of the 5-methylmercaptothiophene-2-carboxaldehyde, and 3 drops of piperidine in 2 mL of ethanol was stirred at 90°C for 3 h. After cooling, the precipitate was filtered, washed with cold ethanol, and dried to yield 246.6 mg (90%) of the title compound as a orange solid.

5.14. Synthesis of 3-[(2-Methylthien-5-yl)methylene]-2-indolinone (SU5424)



A reaction mixture of 134.0 mg of xindole, 151.42 mg of the 5-methylthiophene-2-carboxaldehyde, and 3 drops of piperidine in 2 mL of ethanol was stirred at 90°C for 3 h. After cooling, the precipitate was filtered, washed with cold 5 ethanol, and dried to yield 237.8 mg (99%) of the title compound as a yellow solid.

5.15. Synthesis of 3-[(3-Methylthien-2-yl)methylene]-2-indolinone (SU5427)

10 A reaction mixture of 134.0 mg of oxindole, 151.4 mg of the 3-methylthiophene-2-carboxaldehyde, and 3 drops of piperidine in 2 mL of ethanol was stirred at 90°C for 3 h. After cooling, the precipitate was filtered, washed with cold ethanol, and dried to yield 157.8 mg (65%) of the title  
15 compound as a yellow solid.

5.16. Synthesis of 3-(2,5-Dimethoxybenzylidenyl)-2-indolinone (SU4793)

3-(2,5-Dimethoxybenzylidenyl)-2-indolinone is synthesized according to Method A.  
20

5.17. Synthesis of 3-(2,3-dimethoxybenzylidenyl)-2-indolinone (SU4794)

3-(2,3-dimethoxybenzylidenyl)-2-indolinone is synthesized according to Method A.  
25

5.18. Synthesis of 3-(3-bromo-6-methoxybenzylidenyl)-2-indolinone (SU4796)

3-(3-bromo-6-methoxybenzylidenyl)-2-indolinone is synthesized according to Method A.

30

5.19. Synthesis of 3-[4-(4-t-butylcarbonyl-piperazin-1-yl)benzylidenyl]-2-indolinone (SU5393)

3-[4-(4-t-butylcarbonyl-piperazin-1-yl)benzylidenyl]-2-indolinone is synthesized according to Method B.

35

5.20. Synthesis of 3-[(furan-2-yl)methylene]-2-indolinone (SU4798)

3-[(furan-2-yl)methylene]-2-indolinone is synthesized according to Method A.

- 5.21.        Synthesis of 3-(4-acetamidobenzylidenyl)-2-indolinone (SU4799)  
5        3-(4-acetamidobenzylidenyl)-2-indolinone is synthesized according to Method A.
- 5.22.        Synthesis of 3-(2-chloro-4-hydroxybenzylidenyl)-2-indolinone (SU4932)  
10        3-(2-chloro-4-hydroxybenzylidenyl)-2-indolinone is synthesized according to Method A.
- 5.23.        Synthesis of 3-(4-Bromobenzylidenyl)-2-indolinone (SU4942)  
15        3-(4-Bromobenzylidenyl)-2-indolinone is synthesized according to Method A.
- 5.24.        Synthesis of 3-(4-Acetylaminobenzylidenyl)-2-indolinone (SU4944)  
20        3-(4-Acetylaminobenzylidenyl)-2-indolinone is synthesized according to Method A.
- 5.25.        Synthesis of 3-(2-Methoxybenzylidenyl)-2-indolinone (SU4949)  
25        3-(2-Methoxybenzylidenyl)-2-indolinone is synthesized according to Method A.
- 5.26.        Synthesis of 3-(4-Dimethylaminobenzylidenyl)-1-methyl-2-indolinone (SU4952)  
30        3-(4-Dimethylaminobenzylidenyl)-1-methyl-2-indolinone is synthesized according to Method A.
- 5.27.        Synthesis of 3-(4-Dimethylaminobenzylidenyl)-2-indolinone (SU4312)  
35        3-(4-Dimethylaminobenzylidenyl)-2-indolinone is available from Maybridge Chemical Co. Ltd.

- 5.28. Synthesis of 3-(4-Bromobenzylidenyl)-1-methyl-2-indolinone (SU4956)  
3-(4-Bromobenzylidenyl)-1-methyl-2-indolinone is synthesized according to Method A.
- 5  
5.29. Synthesis of 5-Chloro-3-(4-dimethylaminobenzylidenyl)-2-indolinone (SU4967)  
5-Chloro-3-(4-dimethylaminobenzylidenyl)-2-indolinone is synthesized according to Method A.
- 10  
5.30. Synthesis of 3-(4-Bromobenzylidenyl)-5-chloro-2-indolinone (SU4972)  
3-(4-Bromobenzylidenyl)-5-chloro-2-indolinone is synthesized according to Method A.
- 15  
5.31. Synthesis of 3-(4-Diethylaminobenzylidenyl)-2-indolinone (SU4978)  
3-(4-Diethylaminobenzylidenyl)-2-indolinone is synthesized according to Method A.
- 20  
5.32. Synthesis of 3-(4-Di-n-butylaminobenzylidenyl)-2-indolinone (SU4979)  
3-(4-Di-n-butylaminobenzylidenyl)-2-indolinone is synthesized according to Method A.
- 25  
5.33. Synthesis of 1-Methyl-3-[4-(morpholin-4-yl)benzylidenyl]-2-indolinone (SU4982)  
1-Methyl-3-[4-(morpholin-4-yl)benzylidenyl]-2-indolinone is synthesized according to Method B.
- 30  
5.34. Synthesis of 5-Chloro-3-(4-(morpholine-4-yl)benzylidenyl)-2-indolinone (SU4983)  
5-Chloro-3-(4-(morpholine-4-yl)benzylidenyl)-2-indolinone is synthesized according to Method B.
- 35  
5.35. Synthesis of 3-(3,4-Dichlorobenzylidenyl)-2-indolinone (SU5201)  
3-(3,4-Dichlorobenzylidenyl)-2-indolinone is synthesized according to Method A.

5.36. Synthesis of 3-(2-Ethoxybenzylidenyl)-2-indolinone (SU5204)

3-(2-Ethoxybenzylidenyl)-2-indolinone is synthesized according to Method A.

5

5.37. Synthesis of 3-(4-Fluorobenzylidenyl)-2-indolinone (SU5205)

3-(4-Fluorobenzylidenyl)-2-indolinone is synthesized according to Method A.

10

5.38. Synthesis of 3-[(Thien-2-yl)methylene]-2-indolinone (SU5208)

3-[(Thien-2-yl)methylene]-2-indolinone is synthesized according to Method A.

15

5.39. Synthesis of 3-(2-Methoxybenzylidenyl)-2-indolinone (SU5214)

3-(2-Methoxybenzylidenyl)-2-indolinone is synthesized according to Method A.

20

5.40. Synthesis of 3-[2-[3,5-Di-(trifluoromethyl)phenyl]furan-5-yl)methylene]-2-indolinone (SU5217)

3-[2-[3,5-Di-(trifluoromethyl)phenyl]furan-5-yl)methylene]-2-indolinone is synthesized according to Method A.

25

5.41. Synthesis of 2,6-Di-(dimethylamino)-3,5-di-[(indolin-2-one-3-ylidenyl)methyl]-phenylcyanide (SU5218)

2,6-Di-(dimethylamino)-3,5-di-[(indolin-2-one-3-ylidenyl)methyl]-phenylcyanide is synthesized according to Method A.

30

5.42. Synthesis of 3-[(3-(2-carboxyethyl)-4-methylpyrrol-5-yl)methylene]-2-indolinone (SU5402)

3-[(3-(2-carboxyethyl)-4-methylpyrrol-5-yl)methylene]-2-indolinone is synthesized according to Method A.

35

- 5.43. Synthesis of 3-[(3,4-Dibrom -5-methylpyrrol-2-yl)methylene]-2-indolinone (SU5403)  
3-[(3,4-Dibromo-5-methylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method B.
- 5  
5.44. Synthesis of 3-[(3,4-Dimethyl-2-formylpyrrole-5-yl)methylene]-2-indolinone (SU5405)  
3-[(3,4-Dimethyl-2-formylpyrrole-5-yl)methylene]-2-indolinone is synthesized according to Method A.
- 10  
5.45. Synthesis of 3-[[4-(2-methoxycarbonyl)ethyl]-3-methylpyrrol-5-yl)methylene]-2-indolinone (SU5407)  
3-[[4-(2-methoxycarbonyl)ethyl]-3-methylpyrrol-5-yl)methylene]-2-indolinone is synthesized according to Method A.
- 15  
5.46. Synthesis of 3-[2-Iodofuran-5-yl)methylene]-2-indolinone (SU5409)  
3-[2-Iodofuran-5-yl)methylene]-2-indolinone is synthesized according to Method A.
- 20  
5.47. Synthesis of 3-[(3-Ethoxycarbonyl-2-methylfuran-5-yl)methylene]-2-indolinone (SU5410)  
3-[(3-Ethoxycarbonyl-2-methylfuran-5-yl)methylene]-2-indolinone is synthesized according to Method A.
- 25  
5.48. Synthesis of 3-[(3-Bromothiophene-2-yl)methylene]-2-indolinone (SU5418)  
3-[(3-Bromothiophene-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 30  
5.49. Synthesis of 3-[(2-Chlorothiophene-5-yl)methylene]-2-indolinone (SU5420)  
3-[(2-Chlorothiophene-5-yl)methylene]-2-indolinone is synthesized according to Method A.
- 35  
5.50. Synthesis of 3-[(2,3-Dimethylfuran-5-yl)methylene]-2-indolinone (SU5421)

3-[(2,3-Dimethylfuran-5-yl)methylene]-2-indolinone is synthesized according to Method A.

5.51.        Synthesis of 3-[(5-Nitrothien-2-yl)methylene]-  
5                2-indolinone (SU5422)  
3-[(5-Nitrothien-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.52.        Synthesis of 3-[(2-Carboxythien-5-  
10                yl)methylene]-2-indolinone (SU5423)  
3-[(2-Carboxythien-5-yl)methylene]-2-indolinone is synthesized according to Method A.

5.53.        Synthesis of 3-[(2-Bromothiène-5-  
15                yl)methylene]-2-indolinone (SU5425)  
3-[(2-Bromothiène-5-yl)methylene]-2-indolinone is synthesized according to Method A.

5.54.        Synthesis of 3-[(4-Bromothiène-2-  
20                yl)methylene]-2-indolinone (SU5426)  
3-[(4-Bromothiène-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.55.        Synthesis of 3-[(2-Sulphonylfuran-5-  
25                yl)methylene]-2-indolinone sodium salt  
                 (SU5428)  
3-[(2-Sulphonylfuran-5-yl)methylene]-2-indolinone sodium salt is synthesized according to Method A.

5.56.        Synthesis of 3-[(Furan-2-yl)methylene]-2-  
30                indolinone (SU5429)  
3-[(Furan-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.57.        Synthesis of 3-[(2-Methylfuran-5-  
35                yl)methylene]-2-indolinone (SU5430)  
3-[(2-Methylfuran-5-yl)methylene]-2-indolinone is synthesized according to Method A.

- 5.58.        Synthesis of 3-[(2-Ethylfuran-5-yl)methylene-2-indolinone (SU5431)  
3-[(2-Ethylfuran-5-yl)methylene-2-indolinone is synthesized according to Method A.
- 5        5.59.        Synthesis of 3-[(2-Nitrofuran-5-yl)methylene]-2-indolinone (SU5432)  
3-[(2-Nitrofuran-5-yl)methylene]-2-indolinone is synthesized according to Method A.
- 10       5.60.        Synthesis of 3-[(5-Bromofuran-2-yl)methylene]-2-indolinone (SU5438)  
3-[(5-Bromofuran-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 15       5.61.        Synthesis of 3-[(2-Ethylthien-5-yl)methylene]-2-indolinone (SU5451)  
3-[(2-Ethylthien-5-yl)methylene]-2-indolinone is synthesized according to Method A.
- 20       5.62.        Synthesis of 3-[(4,5-Dimethyl-3-ethylpyrrol-2-yl)methylene]-2-indolinone (SU5453)  
3-[(4,5-Dimethyl-3-ethylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 25       5.63.        Synthesis of 3-[(5-Ethoxycarbonyl-4-ethoxycarbonylethyl-3-ethoxycarbonylmethylpyrrol-2-yl)methylene]-2-indolinone (SU5454)  
3-[(5-Ethoxycarbonyl-4-ethoxycarbonylethyl-3-ethoxycarbonylmethylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 30       5.64.        Synthesis of 3-[(5-Carboxy-3-ethyl-4-methylpyrrol-2-yl)methylene]-2-indolinone (SU5455)  
3-[(5-Carboxy-3-ethyl-4-methylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 35

- 5.65. Synthesis of 3-[(3,5-Diiodo-4-methylpyrrol-2-yl)methylene]-2-indolinone (SU5456)  
3-[(3,5-Diiodo-4-methylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 5 5.66. Synthesis of 3-[(5-Chloro-3-methoxycarbonyl-4-methoxycarbonylmethylpyrrol-2-yl)methylene]-2-indolinone (SU5459)  
3-[(5-Chloro-3-methoxycarbonyl-4-methoxycarbonylmethylpyrrol-2-yl)methylene]-2-indolinone is  
10 synthesized according to Method A.
- 5.67. Synthesis of 3-[(3-Acetyl-5-ethoxycarbonyl-4-methylpyrrol-2-yl)methylene]-2-indolinone (SU5460)  
3-[(3-Acetyl-5-ethoxycarbonyl-4-methylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to  
15 Method A.
- 5.68. Synthesis of 3-[[1-(3,5-Dichlorophenyl)pyrrol-2-yl]methylene]-2-indolinone (SU5461)  
20 3-[[1-(3,5-Dichlorophenyl)pyrrol-2-yl]methylene]-2-indolinone is synthesized according to Method A.
- 5.69. Synthesis of 3-[1-(4-Chlorophenyl)pyrrol-2-yl)methylene]-2-indolinone (SU5462)  
25 3-[1-(4-Chlorophenyl)pyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 5.70. Synthesis of 3-[(4-Ethoxycarbonyl-3-methylpyrrol-2-yl)methylene]-2-indolinone (SU5463)  
30 3-[(4-Ethoxycarbonyl-3-methylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 5.71. Synthesis of 3-[(1-Methylpyrrol-2-yl)methylene]-2-indolinone (SU5464)  
35 3-[(1-Methylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.



5.72.        Synthesis of 3-[(5-Ethoxycarbonyl-3-ethoxycarbonylethyl-4-ethoxycarbonylmethylpyrrol-2-yl)methylene]-2-indolinone (SU5465)

3-[(5-Ethoxycarbonyl-3-ethoxycarbonylethyl-4-ethoxycarbonylmethylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.73.        Synthesis of 3-[4-(Pyrrolidin-1-yl)benzylidenyl]-2-indolinone (SU5466)

10        3-[4-(Pyrrolidin-1-yl)benzylidenyl]-2-indolinone is synthesized according to Method A.

5.74.        Synthesis of 3-[(5-Methylimidazol-2-yl)methylene]-2-indolinone (SU5468)

15        3-[(5-Methylimidazol-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.75.        Synthesis of 3-[(5-Methylthiazol-2-yl)methylene]-2-indolinone (SU5469)

20        3-[(5-Methylthiazol-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.76.        Synthesis of 3-[(3-Methylpyrazol-5-yl)methylene]-2-indolinone (SU5472)

25        3-[(3-Methylpyrazol-5-yl)methylene]-2-indolinone is synthesized according to Method A.

5.77.        Synthesis of 3-[(Imidazol-4-yl)methylene]-2-indolinone (SU5473)

30        3-[(Imidazol-4-yl)methylene]-2-indolinone is synthesized according to Method A.

5.78.        Synthesis of 3-[(4-Chloropyrazol-3-yl)methylene]-2-indolinone (SU5474)

35        3-[(4-Chloropyrazol-3-yl)methylene]-2-indolinone is synthesized according to Method A.

- 5.79. Synthesis of 3-[(4-Bromo-1-(4-chlorobenzyl)pyrazol-5-yl)methylene]-2-indolinone (SU5475)  
3-[(4-Bromo-1-(4-chlorobenzyl)pyrazol-5-yl)methylene]-2-indolinone is synthesized according to Method A.  
5
- 5.80. Synthesis of 3-[(4-Chloro-1-methylpyrazol-3-yl)methylene]-2-indolinone (SU5476)  
3-[(4-Chloro-1-methylpyrazol-3-yl)methylene]-2-indolinone is synthesized according to Method A.  
10
- 5.81. Synthesis of 3-[(4-Ethyl-3,5-dimethylpyrrol-2-yl)methylene]-2-indolinone (SU5477)  
3-[(4-Ethyl-3,5-dimethylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method B.  
15
- 5.82. Synthesis of 3-[(5-Ethylpyrrol-2-yl)methylene]-2-indolinone (SU5478)  
3-[(5-Ethylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method B.  
20
- 5.83. Synthesis of 3-[3,5-Dimethyl-4-(propen-2-yl)pyrrol-2-yl)methylene]-2-indolinone (SU5479)  
3-[3,5-Dimethyl-4-(propen-2-yl)pyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method B.  
25
- 5.84. Synthesis of 5,6-Dimethoxyl-3-[2,3-dimethoxylbenzylidenyl]-2-indolinone (SU5495)  
5,6-Dimethoxyl-3-[2,3-dimethoxylbenzylidenyl]-2-indolinone is synthesized according to Method A.  
30
- 5.85. Synthesis of 3-[2,4,6-Trimethoxybenzylidenyl]-2-indolinone (SU5607)  
3-[2,4,6-Trimethoxybenzylidenyl]-2-indolinone is synthesized according to Method A.  
35
- 5.86. Synthesis of 5-Chloro-3-[(pyrrol-2-yl)methylene]-2-indolinone (SU5612)

5-Chloro-3-[(pyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.87. Synthesis of 5-Chloro-3-[(3-methylpyrrol-2-yl)methylene]-2-indolinone (SU5613)  
5 5-Chloro-3-[(3-methylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.88. Synthesis of 3-(4-isopropylbenzylidenyl)-2-indolinone (SU4313)  
10 3-(4-isopropylbenzylidenyl)-2-indolinone is available from Maybridge Chemical Co. Ltd.

5.89. Synthesis of 5-Chloro-3-[(3,5-dimethylpyrrol-2-yl)methylene]-2-indolinone (SU5614)  
15 5-Chloro-3-[(3,5-dimethylpyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.90. Synthesis of 3-[(pyrrol-2-yl)methylene]-2-indolinone (SU4314)  
20 3-[(pyrrol-2-yl)methylene]-2-indolinone is available from Maybridge Chemical Co. Ltd.

5.91. Synthesis of 5-Chloro-3-[(indol-3-yl)methylene]-2-indolinone (SU5615)  
25 5-Chloro-3-[(indol-3-yl)methylene]-2-indolinone is synthesized according to Method A.

5.92. Synthesis of 5-Chloro-3-[(thien-2-yl)methylene]-2-indolinone (SU5616)  
30 5-Chloro-3-[(thien-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.93. Synthesis of 5-Chloro-3-[(3-methylthien-2-yl)methylene]-2-indolinone (SU5617)  
35 5-Chloro-3-[(3-methylthien-2-yl)methylene]-2-indolinone is synthesized according to Method A.

- 5.94. Synthesis of 5-Chloro-3-[(5-methylthien-2-yl)methylene]-2-indolinone (SU5618)  
5-Chloro-3-[(5-methylthien-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 5 5.95. Synthesis of 5-Chloro-3-[(5-ethylthien-2-yl)methylene]-2-indolinone (SU5619)  
5-Chloro-3-[(5-ethylthien-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 10 5.96. Synthesis of 5-Chloro-3-[(5-methylmercaptothien-2-yl)methylene]-2-indolinone (SU5620)  
5-Chloro-3-[(5-methylmercaptothien-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 15 5.97. Synthesis of 5-Chloro-3-[(imidazol-2-yl)methylene]-2-indolinone (SU5621)  
5-Chloro-3-[(imidazol-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 20 5.98. Synthesis of 3-[2,4-Dimethoxy-6-methylbenzylidenyl]-2-indolinone (SU5623)  
3-[2,4-Dimethoxy-6-methylbenzylidenyl]-2-indolinone is synthesized according to Method A.
- 25 5.99. Synthesis of 5-Nitro-3-[(pyrrol-2-yl)methylene]-2-indolinone (SU5624)  
5-Nitro-3-[(pyrrol-2-yl)methylene]-2-indolinone is synthesized according to Method A.
- 30 5.100. Synthesis of 3-[(3-Methylpyrrol-2-yl)methylene]-5-nitro-2-indolinone (SU5625)  
3-[(3-Methylpyrrol-2-yl)methylene]-5-nitro-2-indolinone is synthesized according to Method A.
- 35 5.101. Synthesis of 3-[(3,5-Dimethylpyrrol-2-yl)methylene]-5-nitro-2-indolinone (SU5626)

3-[(3,5-Dimethylpyrrol-2-yl)methylene]-5-nitro-2-indolinone is synthesized according to Method A.

5.102. Synthesis of 3-[(Ind 1-3-yl)methylene]-5-nitro-2-indolinone (SU5627)

3-[(Indol-3-yl)methylene]-5-nitro-2-indolinone is synthesized according to Method A.

5.103. Synthesis of 5-Nitro-3-[(thien-2-yl)methylene]-2-indolinone (SU5628)

5-Nitro-3-[(thien-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.104. Synthesis of 3-[(3-Methylthien-2-yl)methylene]-5-nitro-2-indolinone (SU5629)

3-[(3-Methylthien-2-yl)methylene]-5-nitro-2-indolinone is synthesized according to Method A.

5.105. Synthesis of 3-[(5-Methylthien-2-yl)methylene]-5-nitro-2-indolinone (SU5630)

3-[(5-Methylthien-2-yl)methylene]-5-nitro-2-indolinone is synthesized according to Method A.

5.106. Synthesis of 3-[(5-Ethylthien-2-yl)methylene]-5-nitro-2-indolinone (SU5631)

3-[(5-Ethylthien-2-yl)methylene]-5-nitro-2-indolinone is synthesized according to Method A.

5.107. Synthesis of 3-[(5-Methylmercaptothien-2-yl)methylene]-5-nitro-2-indolinone (SU5632)

3-[(5-Methylmercaptothien-2-yl)methylene]-5-nitro-2-indolinone is synthesized according to Method A.

5.108. Synthesis of 3-[(Imidazol-2-yl)methylene]-5-nitro-2-indolinone (SU5633)

3-[(Imidazol-2-yl)methylene]-5-nitro-2-indolinone is synthesized according to Method A.

5.109. Synthesis of 3-[(Oxazol-2-yl)methylene]-2-indolinone (CS7127)

3-[(Oxazol-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.110. Synthesis of 3-[(Oxazol-4-yl)methylene]-2-indolinone (CS7128)

3-[(Oxazol-4-yl)methylene]-2-indolinone is synthesized according to Method A.

5.111. Synthesis of 3-[(Oxazol-5-yl)methylene]-2-indolinone (CS7129)

3-[(Oxazol-5-yl)methylene]-2-indolinone is synthesized according to Method A.

5.112. Synthesis of 3-[(Thiazol-2-yl)methylene]-2-indolinone (CS7130)

3-[(Thiazol-2-yl)methylene]-2-indolinone is synthesized according to Method A.

5.113. Synthesis of 3-[(Thiazol-4-yl)methylene]-2-indolinone (CS7131)

3-[(Thiazol-4-yl)methylene]-2-indolinone is synthesized according to Method A.

5.114. Synthesis of 3-[(Thiazol-5-yl)methylene]-2-indolinone (CS7132)

3-[(Thiazol-5-yl)methylene]-2-indolinone is synthesized according to Method A.

5.115. Synthesis of 3-[(Imidazol-2-yl)methylene]-2-indolinone (CS7133)

3-[(Imidazol-2-yl)methylene]-2-indolinone is synthesized according to Method A.

- 5.116. Synthesis of 3-[(Pyrazol-3-yl)methylene]-2-indolinone (CS7135)  
3-[(Pyrazol-3-yl)methylene]-2-indolinone is synthesized according to Method A.
- 5 5.117. Synthesis of 3-[(Pyrazol-4-yl)methylene]-2-indolinone (CS7136)  
3-[(Pyrazol-4-yl)methylene]-2-indolinone is synthesized according to Method A.
- 10 5.118. Synthesis of 3-[(Isoxazol-3-yl)methylene]-2-indolinone (CS7137)  
3-[(Isoxazol-3-yl)methylene]-2-indolinone is synthesized according to Method A.
- 15 5.119. Synthesis of 3-[(Isoxazol-4-yl)methylene]-2-indolinone (CS7138)  
3-[(Isoxazol-4-yl)methylene]-2-indolinone is synthesized according to Method A.
- 20 5.120. Synthesis of 3-[(Isoxazol-5-yl)methylene]-2-indolinone (CS7139)  
3-[(Isoxazol-5-yl)methylene]-2-indolinone is synthesized according to Method A.
- 25 5.121. Synthesis of 3-[(Isothiazol-3-yl)methylene]-2-indolinone (CS7140)  
3-[(Isothiazol-3-yl)methylene]-2-indolinone is synthesized according to Method A.
- 30 5.122. Synthesis of 3-[(Isothiazol-4-yl)methylene]-2-indolinone (CS7141)  
3-[(Isothiazol-4-yl)methylene]-2-indolinone is synthesized according to Method A.
- 35 5.123. Synthesis of 3-[(Isothiazol-5-yl)methylene]-2-indolinone (CS7142)  
3-[(Isothiazol-5-yl)methylene]-2-indolinone is synthesized according to Method A.

5.124. Synthesis of 3-[(1,2,3-Triazol-4-yl)methylene]-2-indolinone (CS7143)  
3-[(1,2,3-Triazol-4-yl)methylene]-2-indolinone is synthesized according to Method A.

5 5.125. Synthesis of 3-[(1,3,4-Thiadiazol-2-yl)methylene]-2-indolinone (CS7144)  
3-[(1,3,4-Thiadiazol-2-yl)methylene]-2-indolinone is synthesized according to Method A.

10 5.126. Synthesis of 3-[(5-Phenyl-1,2,4-oxadiazol-3-yl)methylene]-2-indolinone (CS7145)  
3-[(5-Phenyl-1,2,4-oxadiazol-3-yl)methylene]-2-indolinone is synthesized according to Method A.

15 5.127. Synthesis of 3-[(3-Phenyl-1,2,4-oxadiazol-5-yl)methylene]-2-indolinone (CS7146)  
3-[(3-Phenyl-1,2,4-oxadiazol-5-yl)methylene]-2-indolinone is synthesized according to Method A.

20 5.128. Synthesis of 3-[(3-Phenyl-1,2,5-oxadiazol-4-yl)methylene]-2-indolinone (CS7147)  
3-[(3-Phenyl-1,2,5-oxadiazol-4-yl)methylene]-2-indolinone is synthesized according to Method A.

## 6. EXAMPLES: In Vitro RTK Assays

25 The following in vitro assays may be used to determine the level of activity and effect of the different compounds of the present invention on one or more of the RTKs. Similar assays can be designed along the same lines for any tyrosine kinase using techniques  
30 well known in the art.

### 6.1. Enzyme Linked Immunosorbent Assay (ELISA)

Enzyme linked immunosorbent assays (ELISA) may be used to detect and measure the presence of tyrosine  
35 kinase activity. The ELISA may be conducted according to known protocols which are described in, for example,



Voller, et al., 1980, "Enzyme-Linked Immunosorbent Assay," In: *Manual of Clinical Immunology*, 2d ed., edited by Rose and Friedman, pp 359-371 Am. Soc. Of Microbiology, Washington, D.C.

- 5 The disclosed protocol may be adapted for determining activity with respect to a specific RTK. For example, the preferred protocols for conducting the ELISA experiments for specific RTKs is provided below. Adaptation of these protocols for determining a  
10 compound's activity for other members of the RTK family, as well as non-receptor tyrosine kinases, are within the scope of those in the art.

#### 6.1.1. FLK-1 ELISA

- 15 An ELISA assay was conducted to measure the kinase activity of the FLK-1 receptor and more specifically, the inhibition or activation of protein tyrosine kinase activity on the FLK-1 receptor. Specifically, the following assay was conducted to  
20 measure kinase activity of the FLK-1 receptor in FLK-1/NIH3T3 cells.

##### Materials And Methods.

Materials. The following reagents and supplies were used:

- 25 a. Corning 96-well ELISA plates (Corning Catalog No. 25805-96);  
b. Cappel goat anti-rabbit IgG (catalog no. 55641);  
c. PBS (Gibco Catalog No. 450-1300EB);  
30 d. TBSW Buffer (50 mM Tris (pH 7.2), 150 mM NaCl and 0.1% Tween-20);  
e. Ethanolamine stock (10% ethanolamine (pH 7.0), stored at 4°C);  
f. HNTG buffer (20mM HEPES buffer (pH 7.5), 150mM  
35 NaCl, 0.2% Triton X-100, and 10% glycerol);  
g. EDTA (0.5 M (pH 7.0) as a 100X stock);

- h. Sodium ortho vanadate (0.5 M as a 100X stock);
- i. Sodium pyro phosphate (0.2M as a 100X stock);
- j. NUNC 96 well V bottom polypropylene plates  
(Applied Scientific Catalog No. AS-72092);
- 5 k. NIH3T3 C7#3 Cells (FLK-1 expressing cells);
- l. DMEM with 1X high glucose L Glutamine (catalog  
No. 11965-050);
- 10 m. FBS, Gibco (catalog no. 16000-028);
- n. L-glutamine, Gibco (catalog no. 25030-016);
- o. VEGF, PeproTech, Inc. (catalog no. 100-20) (kept  
as 1  $\mu$ g/100  $\mu$ l stock in Milli-Q dH<sub>2</sub>O and stored  
at -20°C;
- 15 p. Affinity purified anti-FLK-1 antiserum,  
Enzymology Lab, Sugen, Inc.;
- q. UB40 monoclonal antibody specific for  
phosphotyrosine, Enzymology Lab, Sugen, Inc.  
(see, Fendley, et al., 1990, *Cancer Research*  
50:1550-1558);
- 20 r. EIA grade Goat anti-mouse IgG-POD (BioRad  
catalog no. 172-1011);
- s. 2,2-azino-bis(3-ethylbenz-thiazoline-6-sulfonic  
acid (ABTS) solution (100mM citric acid  
(anhydrous), 250 mM Na<sub>2</sub>HPO<sub>4</sub> (pH 4.0), 0.5 mg/ml  
ABTS (Sigma catalog no. A-1888)), solution  
25 should be stored in dark at 4°C until ready for  
use;
- t. H<sub>2</sub>O<sub>2</sub> (30% solution) (Fisher catalog no. H325);
- u. ABTS/H<sub>2</sub>O<sub>2</sub> (15ml ABTS solution, 2  $\mu$ l H<sub>2</sub>O<sub>2</sub>)  
prepared 5 minutes before use and left at room  
temperature;
- 30 v. 0.2 M HCl stock in H<sub>2</sub>O;
- w. dimethylsulfoxide (100%) (Sigma Catalog No. D-  
8418); and
- Y. Trypsin-EDTA (Gibco BRL Catalog No. 25200-049).
- 35 Protocol. The following protocol was used for  
conducting the assay:

1. Coat Corning 96-well elisa plates with 1.0µg per well Cappel Anti-rabbit IgG antibody in 0.1M Na<sub>2</sub>CO<sub>3</sub> pH 9.6. Bring final volume to 150 µl per well. Coat plates overnight at 4°C. Plates can be kept up to two weeks 5 when stored at 4°C.
2. Grow cells in Growth media(DMEM, supplemental with 2.0mM L-Glutamine, 10% FBS) in suitable culture dishes until confluent at 37°C, 5% CO<sub>2</sub>.
3. Harvest cells by trypsinization and seed in 10 Corning 25850 polystyrene 96-well roundbottom cell plates, 25,000 cells/well in 200µl of growth media.
4. Grow cells at least one day at 37°C, 5% CO<sub>2</sub>.
5. Wash cells with D-PBS 1X.
6. Add 200µl/well of starvation media (DMEM, 2.0mM 15 l-Glutamine, 0.1% FBS). Incubate overnight at 37°C, 5% CO<sub>2</sub>.
7. Dilute Compounds/Extracts 1:20 in polypropylene 96 well plates using starvation media. Dilute dimethylsulfoxide 1:20 for use in control wells.
- 20 8. Remove starvation media from 96 well cell culture plates and add 162 µl of fresh starvation media to each well.
9. Add 18µl of 1:20 diluted Compound/Extract dilution (from step 7) to each well plus the 1:20 25 dimethylsulfoxide dilution to the control wells (+/- VEGF), for a final dilution of 1:200 after cell stimulation. Final dimethylsulfoxide is 0.5 %. Incubate the plate at 37°C, 5% CO<sub>2</sub> for two hours.
10. Remove unbound antibody from ELISA plates by 30 inverting plate to remove liquid. Wash 3 times with TBSW + 0.5% ethanolamine, pH 7.0. Pat the plate on a paper towel to remove excess liquid and bubbles.
11. Block plates with TBSW + 0.5% Ethanolamine, pH 7.0, 150 µl per well. Incubate plate thirty minutes 35 while shaking on a microtiter plate shaker.
12. Wash plate 3 times as described in step 10.

13. Add 0.5 $\mu$ g/well affinity purified anti-FLU-1 polyclonal rabbit antiserum. Bring final volume to 150 $\mu$ l/well with TBSW + 0.5% ethanolamine pH 7.0. Incubate plate for thirty minutes while shaking.
- 5 14. Add 180  $\mu$ l starvation medium to the cells and stimulate cells with 20 $\mu$ l/well 10.0mM sodium ortho vanadate and 500 ng/ml VEGF (resulting in a final concentration of 1.0mM sodium ortho vanadate and 50ng/ml VEGF per well) for eight minutes at 37°C, 5% CO<sub>2</sub>.
- 10 Negative control wells receive only starvation medium.
15. After eight minutes, media should be removed from the cells and washed one time with 200 $\mu$ l/well PBS.
16. Lyse cells in 150 $\mu$ l/well HNTG while shaking at room temperature for five minutes. HNTG formulation includes sodium ortho vanadate, sodium pyro phosphate and EDTA.
17. Wash ELISA plate three times as described in step 10.
18. Transfer cell lysates from the cell plate to 20 elisa plate and incubate while shaking for two hours. To transfer cell lysate pipette up and down while scrapping the wells.
19. Wash plate three times as described in step 10.
20. Incubate ELISA plate with 0.02 $\mu$ g/well UB40 in 25 TBSW + 0.5% ethanolamine. Bring final volume to 150 $\mu$ l/well. Incubate while shaking for 30 minutes.
21. Wash plate three times as described in step 10.
22. Incubate ELISA plate with 1:10,000 diluted EIA grade goat anti-mouse IgG conjugated horseradish 30 peroxidase in TBSW + 0.5% ethanolamine, pH 7.0. Bring final volume to 150 $\mu$ l/well. Incubate while shaking for thirty minutes.
23. Wash plate as described in step 10.
24. Add 100  $\mu$ l of ABTS/H<sub>2</sub>O<sub>2</sub> solution to well.
- 35 Incubate ten minutes while shaking.
25. Add 100  $\mu$ l of 0.2 M HCl for 0.1 M HCl final to stop the color development reaction. Shake 1 minute at

room temperature. Remove bubbles with slow stream of air and read the ELISA plate in an ELISA plate reader at 410 nm.

### 6.1.2. HER-2 ELISA

5 Assay 1: EGF Receptor-HER2 Chimeric Receptor Assay In Whole Cells. HER2 kinase activity in whole EGFR-NIH3T3 cells was measured as described below:

*Materials and Reagents.* The following materials and reagents were used to conduct the assay:

- 10 a. EGF: stock concentration= 16.5 ILM; EGF 201, TOYOBO, Co., Ltd. Japan.
- b. 05-101 (UBI) (a monoclonal antibody recognizing an EGFR extracellular domain).
- 15 c. Anti-phosphotyrosine antibody (anti-Ptyr) (polyclonal) (see, Fendley, et al., supra).
- d. Detection antibody: Goat anti-rabbit IgG horse radish peroxidase conjugate, TAGO, Inc., Burlingame, CA.
- e. TBST buffer:  
20 Tris-HCl, pH 7.2 50 mM  
NaCl 150 mM  
Triton X-100 0.1
- f. HNTG 5X stock:  
25 HEPES 0.1 M  
NaCl 0.75 M  
Glycerol 50%  
Triton X-100 1.0%
- g. ABTS stock:  
30 Citric Acid 100 mM  
Na<sub>2</sub>HPO<sub>4</sub> 250 mM  
HCl, conc. 0.5 pH  
ABTS 0.5mg/ml
- \* (2,2' -azinobis(3-ethylbenzthiazolinesulfonic acid)). Keep solution in dark at 4°C until use.
- 35 h. Stock reagents of:  
EDTA 100 mM pH 7.0  
Na<sub>2</sub>VO<sub>4</sub> 0.5 M

Na<sub>4</sub>(P<sub>2</sub>O<sub>7</sub>) 0.2 M

Procedure. The following protocol was used:

A. Pre-coat ELISA Plate

1. Coat ELISA plates (Corning, 96 well, Cat. #25805-96) with 05-101 antibody at 0.5 g per well in PBS, 100 µl final volume/well, and store overnight at 4°C. Coated plates are good for up to 10 days when stored at 4°C.

2. On day of use, remove coating buffer and replace with 100 µl blocking buffer (5% Carnation Instant Non-Fat Dry Milk in PBS). Incubate the plate, shaking, at room temperature (about 23°C to 25°C) for 30 minutes. Just prior to use, remove blocking buffer and wash plate 4 times with TBST buffer.

15 B. Seeding Cells

1. An NIH3T3 cell line overexpressing a chimeric receptor containing the EGFR extracellular domain and extracellular HER2 kinase domain can be used for this assay.

20 2. Choose dishes having 80-90% confluence for the experiment. Trypsinize cells and stop reaction by adding 10% fetal bovine serum. Suspend cells in DMEM medium (10% CS DMEM medium) and centrifuge once at 1500 rpm, at room temperature for 5 minutes.

25 3. Resuspend cells in seeding medium (DMEM, 0.5% bovine serum), and count the cells using trypan blue. Viability above 90% is acceptable. Seed cells in DMEM medium (0.5% bovine serum) at a density of 10,000 cells per well, 100 µl per well, in a 96 well microtiter plate. Incubate seeded cells in 5% CO<sub>2</sub> at 37°C for about 40 hours.

C. Assay Procedures

1. Check seeded cells for contamination using an inverted microscope. Dilute drug stock (10 mg/ml in 35 DMSO) 1:10 in DMEM medium, then transfer 5 µl to a TBST well for a final drug dilution of 1:200 and a final DMSO

concentration of 1%. Control wells receive DMSO alone. Incubate in 5% CO<sub>2</sub> at 37°C for two hours.

2. Prepare EGF ligand: dilute stock EGF in DMEM so that upon transfer of 10 µl dilute EGF (1:125 dilution), 100 nM final concentration is attained.

3. Prepare fresh HNTG' sufficient for 100 µl per well; and place on ice.

HNTG' (10 ml):

	HNTG stock	2.0 ml
10	milli-Q H <sub>2</sub> O	7.3 ml
	EDTA, 100 mM, pH 7.0	0.5 ml
	Na <sub>3</sub> VO <sub>4</sub> , 0.5 M	0.1 ml
	Na <sub>4</sub> (P <sub>2</sub> O <sub>7</sub> ), 0.2 M	0.1 ml

4. After 120 minutes incubation with drug, 15 add prepared SGF ligand to cells, 10 µl per well, to a final concentration of 100 nM. Control wells receive DMEM alone. Incubate, shaking, at room temperature, for 5 minutes.

5. Remove drug, EGF, and DMEM. Wash cells 20 twice with PBS. Transfer HNTG' to cells, 100 µl per well. Place on ice for 5 minutes. Meanwhile, remove blocking buffer from other ELISA plate and wash with TBST as described above.

6. With a pipette tip securely fitted to a 25 micropipettor, scrape cells from plate and homogenize cell material by repeatedly aspirating and dispensing the HNTG' lysis buffer. Transfer lysate to a coated, blocked, and washed ELISA plate. Incubate shaking at room temperature for one hour.

30 7. Remove lysate and wash 4 times with TBST. Transfer freshly diluted anti-Ptyr antibody to ELISA plate at 100 µl per well. Incubate shaking at room temperature for 30 minutes in the presence of the anti-Ptyr antiserum (1:3000 dilution in TBST).

35 8. Remove the anti-Ptyr antibody and wash 4 times with TBST. Transfer the freshly diluted TAGO anti-rabbit IgG antibody to the ELISA plate at 100 µl per

well. Incubate shaking at room temperature for 30 minutes (anti-rabbit IgG antibody: 1:3000 dilution in TBST).

9. Remove TAGO detection antibody and wash 4 times with TBST. Transfer freshly prepared ABTS/H<sub>2</sub>O<sub>2</sub> solution to ELISA plate, 100 µl per well. Incubate shaking at room temperature for 20 minutes. (ABTS/H<sub>2</sub>O<sub>2</sub> solution: 1.0 µl 30% H<sub>2</sub>O<sub>2</sub> in 10 ml ABTS stock).

10. Stop reaction by adding 50 µl 5N H<sub>2</sub>SO<sub>4</sub> (optional), and determine O.D. at 410 nm.

11. The maximal phosphotyrosine signal is determined by subtracting the value of the negative controls from the positive controls. The percent inhibition of phosphotyrosine content for extract-containing wells is then calculated, after subtraction of the negative controls.

**Assay 2: HER-2-BT474 ELISA.** A second assay may be conducted to measure whole cell HER2 activity. Such assay may be conducted as follows:

**Materials And Reagents.** The following materials and reagents were used:

- a. BT-474 (ATCC HBT20), a human breast tumor cell line which expresses high levels of HER2 kinase.
- b. Growth media comprising RPMI + 10% FBS + GMS-G (Gibco supplement) + glutamine for use in growing BT-474 in an incubator with 5% CO<sub>2</sub> at 37°C.
- c. A monoclonal anti-HER2 antibody.
- d. D-PBS:

KH <sub>2</sub> HPO <sub>4</sub>	0.20 g/l	10 (GIBCO, 310-4190AJ)
K <sub>2</sub> HPO <sub>4</sub>	2.16 g/l	
KCl	0.20 g/l	
NaCl	8.00 g/l	(pH 7.2)
- e. Blocking Buffer: TBST plus 5% Milk (Carnation Instant Non-Fat Dry Milk).
- f. TBST buffer:



Tris-HCl 50 mM  
 NaCl 150 mM (pH 7.2, HCl 10 N)  
 Triton X-100 0.1%

wherein stock solution of TES (10X) is prepared, and Triton X-100 is added to the buffer during dilution.

5

g. HNTG buffer (5x):

HEPES 0.1 M  
 NaCl 750 mM (pH 7.2 (HCl, 1 N)  
 Glycerol 50%  
 Triton X-100 1.0%

10

Stock solution (5x) is prepared and kept in 4°C.

h. EDTA-HCl: 0.5 M pH 7.0 (10 N HCl) as 500X stock.

15

i.  $\text{Na}_2\text{VO}_4$ : 0.5 M as 100X stock is kept at -80°C as aliquots.

j.  $\text{Na}_4(\text{P}_2\text{O}_7)$ : 0.2 M as 100X stock.

k. Polyclonal antiserum anti-phosphotyrosine.

20

l. Goat anti-rabbit IgG, horseradish peroxidase (POD) conjugate (detection antibody), Tago (Cat. No. 4520; Lot No. 1802): Tago, Inc., Burlingame, CA.

m. ABTS solution:

Citric acid 100 mM  
 $\text{Na}_2\text{HPO}_4$  250 mM (pH 4.0, 1 N HCl)  
 ABTS 0.5 mg/ml

25

wherein ABTS is 2,2'-azinobis(3-ethylbenzthiazoline sulfonic acid). For this assay, the ABTS solution should be kept in the dark at 4°C. The solution should be discarded when it turns green.

30

n. Hydrogen peroxide: 30% solution is kept in dark and 4°C.

Procedure. All the following steps are at room temperature and aseptically, unless stated otherwise.

35 All ELISA plate washing is by rinsing with distilled water three times and once with TBST.

A. Cell Seeding

1. Grow BT474 cells in tissue culture dishes (Corning 25020-100) to 80-90% confluence and collect using Trypsin-EDTA (0.25%, GIBCO).
- 5 2. Resuspend the cells in fresh medium and transfer to 96-well tissue culture plates (Corning, 25806-96) at about 25,000-50,000 cells/well (100  $\mu$ l/well). Incubate the cells in 5% CO<sub>2</sub> at 37°C overnight.

B. ELISA Plate Coating and Blocking

- 10 1. Coat the ELISA plate (Corning 25805-96) with anti HER2 antibody at 0.5  $\mu$ g/well in 150  $\mu$ l PBS overnight at 4°C, and seal with parafilm. The antibody coated plates can be used up to 2 weeks, when stored at 4°C.
- 15 2. On the day of use, remove the coating solution, replace with 200  $\mu$ l of Blocking Buffer, shake the plate, and then remove the blocking buffer and wash the plate just before adding lysate.

C. Assay Procedures

- 20 1. TBST the drugs in serum-free condition. Before adding drugs, the old media is replaced with serum-free RPMI (90  $\mu$ l/well)
2. Dilute drug stock (in 100% DMSO) 1:10 with RPMI, and transfer 10  $\mu$ l/well of this solution to the  
25 cells to achieve a final drug DMSO concentration at 1%. Incubate the cells in 5% CO<sub>2</sub> at 37°C.
3. Prepare fresh cell lysis buffer (HNTG\*)

5xHNTG	2 ml
EDTA	0.2 ml
Na <sub>3</sub> VO <sub>4</sub>	0.1 ml
Na <sub>2</sub> P <sub>2</sub> O <sub>7</sub>	0.1 ml
H <sub>2</sub> O	7.3 ml
4. After drug preincubation for two hours remove all the solution from the plate, transfer HNTG\*  
35 (100  $\mu$ l/well) to the cells, and shake for 10 minutes.
5. Use a 12-channel pipette to scrape the cells from the plate, and homogenize the lysate by repeat

aspiration and dispensing. Transfer all the lysate to the ELISA plate and shake for 1 hour.

6. Remove the lysate, wash the plate, add anti-pTyr (1:3,000 with TBST) 100  $\mu$ l/well, and shake for 5 30 minutes.

7. Remove anti-pTyr, wash the plate, add goat anti-rabbit IgG conjugated antibody (1:5,000 with TBST) 100  $\mu$ l/well, and shake for 30 minutes.

8. Remove anti-rabbit IgG antibody, wash the 10 plate, and add fresh ABTS/H<sub>2</sub>O<sub>2</sub> (1.2  $\mu$ l H<sub>2</sub>O<sub>2</sub> to 10 ml ABTS) 100 l/well to the plate to start color development, which usually takes 20 minutes.

9. Measure OD 410 nm, Dynatec MR5000.

#### 15 6.1.3. PDGF-R ELISA

All cell culture media, glutamine, and fetal bovine serum were purchased from Gibco Life Technologies (Grand Island, NY) unless otherwise specified. All cells were grown in a humid atmosphere of 20 90-95% air and 5-10% CO<sub>2</sub> at 37°C. All cell lines were routinely subcultured twice a week and were negative for mycoplasma as determined by the Mycotect method (Gibco).

For ELISA assays, cells (U1242, obtained from Joseph Schlessinger, NYU) were grown to 80-90% confluency in 25 growth medium (MEM with 10% FBS, NEAA, 1 mM NaPyr and 2 mM GLN) and seeded in 96-well tissue culture plates in 0.5% serum at 25,000 to 30,000 cells per well. After overnight incubation in 0.5% serum-containing medium, cells were changed to serum-free medium and treated with 30 test compound for 2 hr in a 5% CO<sub>2</sub>, 37°C incubator. Cells were then stimulated with ligand for 5-10 minute followed by lysis with HNTG (20 mM Hepes, 150 mM NaCl, 10% glycerol, 5 mM EDTA, 5 mM Na<sub>2</sub>VO<sub>4</sub>, 0.2% Triton X-100, and 2 mM NaPyr). Cell lysates (0.5 mg/well in PBS) were 35 transferred to ELISA plates previously coated with receptor-specific antibody and which had been blocked with 5% milk in TBST (50 mM Tris-HCl pH 7.2, 150 mM NaCl

and 0.1% Triton X-100) at room temperature for 30 min. Lysates were incubated with shaking for 1 hour at room temperature. The plates were washed with TBST four times and then incubated with polyclonal anti-phosphotyrosine antibody at room temperature for 30 minutes. Excess anti-phosphotyrosine antibody was removed by rinsing the plate with TBST four times. Goat anti-rabbit IgG antibody was added to the ELISA plate for 30 min at room temperature followed by rinsing with TBST four more times. ABTS (100 mM citric acid, 250 mM Na<sub>2</sub>HPO<sub>4</sub> and 0.5 mg/mL 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid)) plus H<sub>2</sub>O<sub>2</sub> (1.2 mL 30% H<sub>2</sub>O<sub>2</sub> to 10 ml ABTS) was added to the ELISA plates to start color development. Absorbance at 410 nm with a reference wavelength of 630 nm was recorded about 15 to 30 min after ABTS addition.

#### 6.1.4. IGF-I ELISA

The following protocol may be used to measure phosphotyrosine level on IGF-I receptor, which indicates IGF-I receptor tyrosine kinase activity.

**Materials And Reagents.** The following materials and reagents were used:

- a. The cell line used in this assay is 3T3/IGF-1R, a cell line which overexpresses IGF-1 receptor.
- 25 b. NIH3T3/IGF-1R is grown in an incubator with 5% CO<sub>2</sub> at 37°C. The growth media is DMEM + 10% FBS (heat inactivated) + 2mM L-glutamine.
- c. Anti-IGF-1R antibody named 17-69 is used. Antibodies are purified by the Enzymology Lab, SUGEN, Inc.
- 30 d. D-PBS:
 

KH <sub>2</sub> PO <sub>4</sub>	0.20 g/l
K <sub>2</sub> HPO <sub>4</sub>	2.16 g/l
KCl	0.20 g/l
NaCl	8.00 g/l (pH 7.2)
- 35 e. Blocking Buffer: TBST plus 5% Milk (Carnation Instant Non-Fat Dry Milk).
- f. TBST buffer:

Tris-HCl	50 mM
NaCl	150mM (pH 7.2/HCl 10N)
Triton X-100	0.1%

5 Stock solution of TBS (10X) is prepared, and Triton X-100 is added to the buffer during dilution.

g. HNTG buffer:

HEPES	20 mM
NaCl	150 mM (pH 7.2/HCl 1N)
Glycerol	10%
Triton X-100	0.2%

10

Stock solution (5X) is prepared and kept at 4°C.

h. EDTA/HCl: 0.5 M pH 7.0 (NaOH) as 100X stock.

15 i.  $\text{Na}_2\text{VO}_4$ : 0.5 M as 100X stock and aliquots are kept in -80°C.

j.  $\text{Na}_2\text{P}_2\text{O}_7$ : 0.2 M as 100X stock.

k. Insulin-like growth factor-1 from Promega (Cat# G5111).

20 l. Polyclonal antiserum anti-phosphotyrosine: rabbit sera generated by Enzymology Lab., SUGEN Inc.

m. Goat anti-rabbit IgG, POD conjugate (detection antibody), Tago (Cat. No. 4520, Lot No. 1802): Tago, Inc., Burlingame, CA.

25 n. ABTS (2,2'-azinobis(3-ethylbenzthiazolinesulfonic acid)) solution:

Citric acid	100 mM
$\text{Na}_2\text{HPO}_4$	250 mM (pH 4.0/1 N HCl)
ABTS	0.5 mg/ml

30 ABTS solution should be kept in dark and 4°C. The solution should be discarded when it turns green.

o. Hydrogen Peroxide: 30% solution is kept in the dark and at 4°C.

Procedure. All the following steps are conducted at 35 room temperature unless it is specifically indicated.

All ELISA plate washings are performed by rinsing the

plate with tap water three times, followed by one TBST rinse. Pat plate dry with paper towels.

A. Cell Seeding:

1. The cells, grown in tissue culture dish 5 (Corning 25020-100) to 80-90% confluence, are harvested with Trypsin-EDTA (0.25%, 0.5 ml/D-100, GIBCO).

2. Resuspend the cells in fresh DMEM + 10% FBS + 2mM L-Glutamine, and transfer to 96 - well tissue culture plate (Corning, 25806-96) at 20,000 cells/well 10 (100  $\mu$ l/well). Incubate for 1 day then replace medium to serum-free medium (90/ $\mu$ l) and incubate in 5% CO<sub>2</sub> and 37°C overnight.

B. ELISA Plate Coating and Blocking:

1. Coat the ELISA plate (Corning 25805-96) 15 with Anti-IGF-1R Antibody at 0.5  $\mu$ g/well in 100  $\mu$ l PBS at least 2 hours.

2. Remove the coating solution, and replace with 100  $\mu$ l Blocking Buffer, and shake for 30 minutes. Remove the blocking buffer and wash the plate just before 20 adding lysate.

C. Assay Procedures:

1. The drugs are tested in serum-free condition.

2. Dilute drug stock (in 100% DMSO) 1:10 with 25 DMEM in 96-well poly-propylene plate, and transfer 10  $\mu$ l/well of this solution to the cells to achieve final drug dilution 1:100, and final DMSO concentration of 1.0%. Incubate the cells in 5% CO<sub>2</sub> at 37°C for 2 hours.

3. Prepare fresh cell lysis buffer (HNTG\*)

30	HNTG	2 ml
	EDTA	0.1 ml
	Na <sub>3</sub> VO <sub>4</sub>	0.1 ml
	Na <sub>4</sub> (P <sub>2</sub> O <sub>7</sub> )	0.1 ml
	H <sub>2</sub> O	7.3 ml

35 4. After drug incubation for two hours, transfer 10  $\mu$ l/well of 200nM IGF-1 Ligand in PBS to the

cells (Final Conc. = 20 nM), and incubate at 5% CO<sub>2</sub> at 37°C for 10 minutes.

5. Remove media and add 100 µl/well HNTG\* and shake for 10 minutes. Look at cells under microscope to see if they are adequately lysed.

6. Use a 12-channel pipette to scrape the cells from the plate, and homogenize the lysate by repeat aspiration and dispense. Transfer all the lysate to the antibody coated ELISA plate, and shake for 1 hour.

10 7. Remove the lysate, wash the plate, transfer anti-pTyr (1:3,000 with TBST) 100 µl/well, and shake for 30 minutes.

8. Remove anti-pTyr, wash the plate, transfer Tago (1:3,000 with TBST) 100 µl/well, and shake for 30 15 minutes.

9. Remove detection antibody, wash the plate, and transfer fresh ABTS/H<sub>2</sub>O<sub>2</sub> (1.2 µl H<sub>2</sub>O<sub>2</sub> to 10 ml ABTS) 100 µl/well to the plate to start color development.

10. Measure OD in Dynatec MR5000, which is 20 connected to Ingres.

#### 6.1.5. EGF Receptor ELISA

EGF Receptor kinase activity (EGFR-NIH3T3 assay) in whole cells was measured as described 25 below:

**Materials and Reagents.** The following materials and reagents were used

- a. EGF Ligand: stock concentration = 16.5 µM; EGF 201, TOYOBO, Co., Ltd. Japan.
- 30 b. 05-101 (UBI) (a monoclonal antibody recognizing an EGFR extracellular domain).
- c. Anti-phosphotyrosine antibody (anti-Ptyr) (polyclonal).
- 35 d. Detection antibody: Goat anti-rabbit IgG horse radish peroxidase conjugate, TAGO, Inc., Burlingame, CA.
- e. TBST buffer:

Tris-HCl, pH 7	50 mM
NaCl	150 mM
Triton X-100	0.1

## f. HNTG 5X stock:

5	HEPES	0.1 M
	NaCl	0.75 M
	Glycerol	50
	Triton X-100	1.0%

## g. ABTS stock:

10	Citric Acid	100 mM
	Na <sub>2</sub> HPO <sub>4</sub>	250 mM
	HCl, conc.	4.0 pH
	ABTS	0.5 mg/ml

Keep solution in dark at 4°C until used.

## h. Stock reagents of:

15	EDTA 100 mM pH 7.0
	Na <sub>2</sub> VO <sub>4</sub> 0.5 M
	Na <sub>4</sub> (P <sub>2</sub> O <sub>7</sub> ) 0.2 M

Procedure. The following protocol was used:

A. Pre-coat ELISA Plate

- 20 1. Coat ELISA plates (Corning, 96 well, Cat. #25805-96) with 05-101 antibody at 0.5 µg per well in PBS, 150 µl final volume/well, and store overnight at 4°C. Coated plates are good for up to 10 days when stored at 4°C.
- 25 2. On day of use, remove coating buffer and replace with blocking buffer (5% Carnation Instant NonFat Dry Milk in PBS). Incubate the plate, shaking, at room temperature (about 23°C to 25°C) for 30 minutes. Just prior to use, remove blocking buffer and wash plate 4 30 times with TBST buffer.

B. Seeding Cells

1. NIH 3T3/C7 cell line (Honegger, et al., Cell 51:199-209, 1987) can be use for this assay.
2. Choose dishes having 80-90% confluence for 35 the experiment. Trypsinize cells and stop reaction by adding 10% CS DMEM medium. Suspend cells in DMEM medium



(10% CS DMEM medium) and centrifuge once at 1000 rpm, and once at room temperature for 5 minutes.

3. Resuspend cells in seeding medium (DMEM, 0.5% bovine serum), and count the cells using trypan blue. Viability above 90% is acceptable. Seed cells in DMEM medium (0.5% bovine serum) at a density of 10,000 cells per well, 100  $\mu$ l per well, in a 96 well microtiter plate. Incubate seeded cells in 5% CO<sub>2</sub> at 37°C for about 40 hours.

10 C. Assay Procedures.

1. Check seeded cells for contamination using an inverted microscope. Dilute drug stock (10 mg/ml in DMSO) 1:10 in DMEM medium, then transfer 5  $\mu$ l to a test well for a final drug dilution of 1:200 and a final DMSO concentration of 1%. Control wells receive DMSO alone. Incubate in 5% CO<sub>2</sub> at 37°C for one hour.

2. Prepare EGF ligand: dilute stock EGF in DMEM so that upon transfer of 10  $\mu$ l dilute EGF (1:12 dilution), 25 nM final concentration is attained.

20 3. Prepare fresh 10 ml HNTG\* sufficient for 100  $\mu$ l per well wherein HNTG\* comprises: HNTG stock (2.0 ml), milli-Q H<sub>2</sub>O (7.3 ml), EDTA, 100 mM, pH 7.0 (0.5 ml), Na<sub>2</sub>VO<sub>4</sub> 0.5 M (0.1 ml) and Na<sub>2</sub>(P<sub>2</sub>O<sub>7</sub>) 0.2 M (0.1 ml).

4. Place on ice.

25 5. After two hours incubation with drug, add prepared EGF ligand to cells, 10  $\mu$ l per well, to yield a final concentration of 25 nM. Control wells receive DMEM alone. Incubate, shaking, at room temperature, for 5 minutes.

30 6. Remove drug, EGF, and DMEM. Wash cells twice with PBS. Transfer HNTG\* to cells, 100  $\mu$ l per well. Place on ice for 5 minutes. Meanwhile, remove blocking buffer from other ELISA plate and wash with TBST as described above.

35 7. With a pipette tip securely fitted to a micropipettor, scrape cells from plate and homogenize cell material by repeatedly aspirating and dispensing the

HNTG' lysis buffer. Transfer lysate to a coated, blocked, and washed ELISA plate. Incubate shaking at room temperature for one hour.

8. Remove lysate and wash 4 times with TBST.  
5 Transfer freshly diluted anti-Ptyr antibody to ELISA plate at 100  $\mu$ l per well. Incubate shaking at room temperature for 30 minutes in the presence of the anti-Ptyr antiserum (1:3000 dilution in TBST).

9. Remove the anti-Ptyr antibody and wash 4  
10 times with TBST. Transfer the freshly diluted TAGO 30 anti-rabbit IgG antibody to the ELISA plate at 100  $\mu$ l per well. Incubate shaking at room temperature for 30 minutes (anti-rabbit IgG antibody: 1:3000 dilution in TBST).

10. Remove detection antibody and wash 4 times  
15 with TBST. Transfer freshly prepared ABTS/H<sub>2</sub>O<sub>2</sub> solution to ELISA plate, 100  $\mu$ l per well. Incubate at room temperature for 20 minutes. ABTS/H<sub>2</sub>O<sub>2</sub> solution: 1.2  $\mu$ l 30% H<sub>2</sub>O<sub>2</sub> in 10 ml ABTS stock.

11. Stop reaction by adding 50  $\mu$ l 5N H<sub>2</sub>SO<sub>4</sub>,  
20 (optional), and determine O.D. at 410 nm.

12. The maximal phosphotyrosine signal is  
determined by subtracting the value of the negative  
controls from the positive controls. The percent  
inhibition of phosphotyrosine content for extract-  
25 containing wells is then calculated, after subtraction of  
the negative controls.

#### 6.1.6. Cellular Insulin Receptor ELISA

The following protocol was used to  
30 determine whether the compounds of the present invention  
possessed insulin receptor tyrosine kinase activity.

**Materials And Reagents.** The following materials and  
reagents were used to measure phosphotyrosine levels on  
the insulin receptor (indicating insulin receptor  
35 tyrosine kinase activity):

1. The preferred cell line was an NIH3T3 cell line (ATCC No. 1658) which overexpresses Insulin Receptor (H25 cells);
2. H25 cells are grown in an incubator with 5% CO<sub>2</sub> at 37°C. The growth media is DMEM + 10% FBS (heat inactivated) + 2mm L-Glutamine;
3. For ELISA plate coating, the monoclonal anti-IR antibody named BBE is used. Said antibodies was purified by the Enzymology Lab, SUGEN, Inc.;
- 10 4. D-PBS, comprising:

KH <sub>2</sub> PO <sub>4</sub>	0.20 g/l (GIBCO, 310-4190AJ)
K <sub>2</sub> HPO <sub>4</sub>	2.16 g/l
KCl	0.20 g/l
NaCl	8.00 g/l (pH 7.2);
- 15 5. Blocking Buffer: TBST plus 5% Milk (Carnation Instant Non-Fat Dry Milk);
6. TBST buffer, comprising:

Tris-HCl	50mM
NaCl	150mM pH 7.2 (HCl, 1 N)
20 Triton X-100	0.1%

Note: Stock solution of TBS (10X) is prepared, and Triton X-100 is added to the buffer during dilution;
7. HNTG buffer, comprising:

HEPES	20mM
25 NaCl	150mM pH 7.2 (HCl, 1 N)
Glycerol	10%
Triton X-100	0.2%

Note: Stock solution (5X) is prepared and kept at 4°C;
8. EDTA.HCl: 0.5 M pH 7.0 (NaOH) as 100X stock;
- 30 9. Na<sub>3</sub>VO<sub>4</sub>: 0.5 M as 100X stock and aliquots are kept in -80°C;
10. Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub>: 0.2 M as 100X stock;
11. Insulin from GIBCO BRL (Cat# 18125039);
12. Polyclonal antiserum Anti-phosphotyrosine:
- 35 rabbit sera generated by Enzymology Lab., SUGEN Inc.;

13. Detection antibody, preferably goat anti-rabbit IgG, POD conjugate, Tago (Cat. No. 4520: Lot No. 1802): Tago, Inc., Burlingame, CA;

14. ABTS solution, comprising:

5 Citric acid 100 mM  
Na<sub>2</sub>HPO<sub>4</sub> 250 mM pH 4.0 (1 N HCl)  
ABTS 0.5 mg/ml

wherein ABTS is 2,2'-azinobis (3-ethylbenathiazoline sulfonic acid) and stored in the dark at 4°C and 10 discarded when it turns green;

15. Hydrogen Peroxide: 30% solution is kept in the dark and at 40°C.

Protocol. All the following steps are conducted at room temperature unless it is specifically indicated.

15 All ELISA plate washings are performed by rinsing the plate with tap water three times, followed by one TBST rinse. All plates were tapped dry with paper towels prior to use.

A. Cell Seeding:

20 1. The cells were grown in tissue culture dish (10 cm, Corning 25020-100) to 80-90% confluence and harvested with Trypsin-EDTA (0.25%, 0.5 ml/D-100, GIBCO);

2. Resuspend the cells in fresh DMEM + 10% FBS + 2mM L-Glutamine, and transfer to 96 - well tissue 25 culture plate (Corning, 25806-96) at 20,000 cells/well (100 µl/well). The cells are then incubated for 1 day. Following such incubation, 0.01% serum medium (90/µl) replaces the old media and the cells incubate in 5% CO<sub>2</sub> and 37°C overnight.

30 B. ELISA Plate Coating and Blocking:

1. Coat the ELISA plate (Corning 25805-96) with Anti-IR Antibody at 0.5 µg/well in 100 µl PBS at least 2 hours.

2. Remove the coating solution, and replace 35 with 100 u; blocking Buffer, and shake for 30 minutes. Remove the blocking buffer and wash the plate just before adding lysate.

### C. Assay Procedures

1. The drugs are tested in serum-free condition.

2. Dilute drug stock (in 100% DMSO) 1:10 with 5 DMEM in 96-well poly-propylene plate, and transfer 10  $\mu$ l/well of this solution to the cells to achieve final drug dilution 1:100, and final DMSO concentration of 1.0%. Incubate the cells in 5% CO<sub>2</sub> at 37°C for 2 hours.

3. Prepare fresh cells lysis buffer (HNTG\*)

10	HNTG (5x)	2 ml
	EDTA	0.1 ml
	Na <sub>3</sub> VO <sub>4</sub>	0.1 ml
	Na <sub>2</sub> P <sub>2</sub> O <sub>7</sub>	0.1 ml
	H <sub>2</sub> O	7.3 ml
15	HNTG*	10 ml

4. After drug incubation for two hours, transfer 10  $\mu$ l/well of 1  $\mu$ M insulin in PBS to the cells (Final concentration = 100 nM), and incubate at 5% CO<sub>2</sub> at 37°C for 10 minutes.

20 5. Remove media and add 100  $\mu$ l/well HNTG\* and shake for 10 minutes. Look at cells under microscope to see if they are adequately lysed.

6. Using a 12-channel pipette, scrape the cells from the plate, and homogenize the lysate by repeat 25 aspiration and dispense. Transfer all the lysate to the antibody coated ELISA plate, and shake for 1 hour.

7. Remove the lysate, wash the plate, transfer anti-pTyr (1:3,000 with TBST) 100  $\mu$ l/well, and shake for 30 minutes.

30 8. Remove anti-pTyr, wash the plate, transfer Tago (1:3,000 with TBST) 100  $\mu$ l/well, and shake for 30 minutes.

9. Remove detection antibody, wash the plate, and transfer fresh ABTS/H<sub>2</sub>O<sub>2</sub> (1.2  $\mu$ l H<sub>2</sub>O<sub>2</sub> to 10 ml ABTS) 35 100  $\mu$ l/well to the plate to start color development.

10. Measure OD in Dynatec MR5000, which is connected to Ingres. All following steps should follow Ingres instruction.

5                    6.1.7.    Experimental Results From ELISA Assays

The experimental results for various compounds according to the invention using the above-described protocols are set forth at Table 1:

10

15

20

25

30

35

TABLE 1

## ELISA Assay Results

	COMPOUND	PDGFR	FLK-1	EGFR	HER2 Kinase	IGF-1R
		IC50 ( $\mu$ M)	IC50 ( $\mu$ M)	IC50 ( $\mu$ M)	IC50 ( $\mu$ M)	IC50 ( $\mu$ M)
5	SU4312	19.4	0.8			
	SU4313	14.5	18.8	11	16.9	8.0
	SU4314	12	0.39			
	SU4793	87.4	4.2			
	SU4794		11.8			
	SU4798		28.8			
10	SU4799		9			
	SU4932		2.2			
	SU4944		8.5			
	SU4952		22.6			
	SU4956				22.5	
	SU4967	7.9	11.2			
15	SU4979		20.9			
	SU4981	33.1	2.1			
	SU4982		21.6		39.4	
	SU4983		4.1			
	SU4984	5.8	1.6		90.2	
	SU5204		4		51.5	
20	SU5205		9.6			
	SU5208		4.7			
	SU5214		14.8	36.7		
	SU5218		6.4			
	SU5401		2.9		89.8	
	SU5402		0.4			
25	SU5403		1.8			
	SU5404	17	0.24			
	SU5405		23.8			
	SU5406		0.17			
	SU5407	53.7	1.1			
	SU5408		0.07			
30	SU5416	10.8	0.11			
	SU5418		15.4			
	SU5419		2.3			
	SU5421		4.6			
	SU5424		2.4			
	SU5425		51.4			
35	SU5427		4.5		70.6	
	SU5428		8.6			
	SU5430		73.4			
	SU5431		41.2			

COMPOUND	PDGFR IC50 ( $\mu$ M)	FLK-1 IC50 ( $\mu$ M)	EGFR IC50 ( $\mu$ M)	HER2 Kinase IC50 ( $\mu$ M)	IGF-1R IC50 ( $\mu$ M)
SU5432		22.8			
SU5450		4.5		92.6	
SU5451		3.4	44		
SU5453	65.5	0.14			
SU5455		36.2			
SU5463		0.18			
SU5464		20.3			
SU5466	86	1.6			
SU5468	55.9	2.7			
SU5472		8.7			
SU5473	14.2	1.5			
SU5474		7.4			
SU5477		0.15			
SU5480		5.3	39.6	30.4	

15

## 6.2. Cell Growth Assays

The following assays may be conducted to measure the effect of the claimed compounds upon cell growth as a result of the compound's interaction with one or more RTKs. Unless otherwise specified, the following assays may be generally applied to measure the activity of a compound against any particular RTK. To the extent that an assay, set forth below, refers to a specific RTK, one skilled in the art would be able to adapt the disclosed protocol for use to measure the activity of a second RTK.

### 6.2.1. Soft Agar Assay

The soft agar assay may be used to measure the effects of substances on cell growth. Unless otherwise stated the soft agar assays were carried out as follows:

**Material And Reagents.** The following materials and reagents were used:

- a. A water bath set at 39°C and another water bath at 37°C.



- b. 2X assay medium is comprised of 2X Dulbecco's 5Modified Eagle's Medium (DMEM) (Gibco Cat. # CA400-4ANO3) supplemented by the following:
- 20% Fetal Bovine Serum (FBS) 2 mM sodium pyruvate 4 mM glutamine amine; and
  - 20 mM HEPES Non-essential Amino Acids (1:50 from 100x stock).
- c. 1X assay medium made of 1X DMEM supplemented with 10% FBS, 1 mM sodium pyruvate, 2 mM glutamine, 10 mM HEPES, non-essential amino acid (1:100 from 100x stock).
- d. 1.6% SeaPlaque Agarose in autoclave bottle.
- e. Sterile 35 mm Corning plates (FMC Bioproducts Cat. #50102).
- f. Sterile 5 ml glass pipets (individually wrapped).
- g. Sterile 15 ml and 50 ml conical centrifuge tubes.
- h. Pipets and sterile tips.
- i. Sterile microcentrifuge tubes.
- j. Cells in T75 flasks: SKOV-3 (ATCC HTB77).
- k. 0.25% Trypsin solution (Gibco #25200-015).

25 Procedure. The following procedure was used to conduct the soft agar assay:

A. Procedure for making the base layer

1. Have all the media warmed up in the 37°C water bath.
2. To make 1X of assay medium + 0.8% agar: make a 1:2 (vol:vol) dilution of melted agar (cooled to 39°C), with 2X assay medium.
3. Keep all media with agar warm in the 39°C water bath when not in use.
4. Dispense 1 ml of 1X assay medium + 0.8% agar into dishes and gently swirl plate to form a uniform base layer. Bubbles should be avoided.

5. Refrigerate base layers to solidify (about 20 minutes). Base layers can be stored overnight in the refrigerator.

B. Procedure for collecting cells

- 5 1. Take out one flask per cell line from the incubator; aspirate off medium; wash once with PBS and aspirate off; add 3 ml of trypsin solution.
2. After all cells dissociate from the flask, add 3 ml of 1X assay media to inhibit trypsin activity.
- 10 Pipet the cells up and down, then transfer the suspension into a 15ml tube.
3. Determine the concentration of cells using a Coulter counter, and the viability by trypan blue exclusion.
- 15 4. Take out the appropriate volume needed to seed 3300 viable cells per plate and dilute it to 1.5 ml with 1X assay medium.

C. Procedure for making the upper 0.4% agarose layer:

- 20 1. Add TBST compounds at twice the desired final assay concentration; + 1.5 ml of cell suspension in 1X assay medium 10% FBS; + 1.5 ml of 1X assay medium + 0.8% agarose: Total = 3.0 ml 1X media 10% FBS + 0.4% agarose with 3300 viable cells/ml, with and without TBST  
25 compounds.

(Made by 1:2 dilution of 2X media with 1.6% agar 30 for the base layer procedure above.)

2. Plate 1 ml of the Assay Mix onto the 1 ml base layer. The duplicates are plated from the 3 ml  
30 volume.
3. Incubate the dishes for 2-3 weeks in a 100% humidified, 10% CO<sub>2</sub> incubator.
4. Colonies that are 60 microns and larger are scored positive.

35

### 6.2.2. Sulf rhodamine B (SRB) Growth Assays

The SRB assays may be used to measure the effects of substances on cell growth. The assays are carried out as follows:

#### 5 Assay 1: 3T3/E/H+TGF- $\alpha$ (T) Cell Growth SRB Assay

##### Materials:

- 96-well flat bottom sterile plates
- 96-well round bottom sterile plates
- sterile 25 ml or 100 ml reservoir
- 10 pipets, multi-channel pipetman
- sterile pipet tips
- sterile 15 ml and 50 ml tubes

##### Reagents:

- 15 0.4% SRB in 1% acetic acid
- 10 mM Tris base
- 10% TCA
- 1% acetic acid
- sterile DMSO (Sigma)
- 20 compound in DMSO (100 mM or less stock solution)
- 25% Trypsin-EDTA in Cell Dissociation Solution (Sigma)

##### Cell line and growth medium:

- 3T3/E/H+TGF- $\alpha$ (T) (NIH 3T3 clone 7 cells expressing EGF-
- 25 R/HER2 chimera and TGF- $\alpha$ , tumor-derived autocrine loop cells)
- 2% calf serum/DMEM + 2 mM glutamine

##### Protocol:

#### 30 Day 0: Cell Plating:

This part of assay is carried out in a laminar flow hood.

1. Trypsinize cells as usual. Transfer 100  $\mu$ l of cell suspension to 10 ml of isotone. Count cells with 35 the Coulter Counter.

2. Dilute cells in growth medium to 60,000 cells/ml. Transfer 100  $\mu$ l of cells to each well in a 96-well flat bottom plate to give 6000 cells/well.
3. Use half of plate (4 rows) for each compound and 5 quadruplicate wells for each compound concentration, a set of 4 wells for medium control and 4 wells for DMSO control.
4. Gently shake plates to allow for uniform attachment of the cells.
- 10 5. Incubate the plates at 37°C in a 10% CO<sub>2</sub> incubator.

Day 1: Addition of Compound:

This part of assay is carried out in a laminar flow hood.

- 15 1. In 96 well-round bottom plate, add 125  $\mu$ l of growth medium to columns 3 to 11. This plate is used to titrate out the compound, 4 rows per compound.
2. In a sterile 15 ml tube, make a 2X solution of  
20 the highest concentration of compound by adding 8  $\mu$ l of the compound to a total of 2 ml growth medium for a dilution of 1:250. At this dilution, the concentration of DMSO is 0.4% for a 2X solution or 0.2% for 1X solution on the cells. The starting concentration of the  
25 compound is usually 100  $\mu$ M but this concentration may vary depending upon the solubility of the compound.
3. Transfer the 2X starting compound solution to quadruplicate wells in column 12 of the 96-well round bottom plate. Do 1:2 serial dilutions across the plate  
30 from right to left by transferring 125  $\mu$ l from column 12 to column 11, column 11 to 10 and so on. Transfer 100  $\mu$ l of compound dilutions onto 100  $\mu$ l medium on cells in corresponding wells of 96-well flat bottom plate. Total volume per well should be 200  $\mu$ l.
- 35 4. For vehicle control, prepare a 2X solution of DMSO at 0.4% DMSO in growth medium. Transfer 100  $\mu$ l of

the DMSO solution to the appropriate wells of cells. The final concentration of DMSO is 0.2%.

5. For the medium control wells, add 100  $\mu$ l/well of growth medium to the appropriate wells of cells.

5 6. Return the plate to the incubator and incubate for 4 days.

#### Day 5: Development of Assay

This part of assay is carried out on the bench.

10

1. Aspirate or pour off medium. Add 200  $\mu$ l cold 10% TCA to each well to fix cells. Incubate plate for at least 60 min. at 4°C.

2. Discard TCA and rinse wells 5 times with water.

15 Dry plates upside down on paper towels.

3. Stain cells with 100  $\mu$ l/well 0.4% SRB for 10 min.

4. Pour off SRB and rinse wells 5 times with 1% acetic acid. Dry plates completely upside down on paper 20 towels.

5. Solubilize dye with 100  $\mu$ l/well 10 mM Tris base for 5-10 min. on shaker.

6. Read plates on Dynatech ELISA Plate Reader at 570 nm with reference at 630 nm.

25

#### Assay 2: 3T3/EGF-R+TGF-a(T) Cell Growth SRB Assay

Materials and Reagents same as for Assay 1.

Cell line and growth medium:

30 3T3/EGF-R+TGF-a(T) (NIH 3T3 clone 7 cells expressing EGF-R and TGF-a, tumor-derived autocrine loop cells)  
2% calf serum/DMEM + 2 mM glutamine

#### Protocol:

35 Day 0: Cell Plating:

This part of assay is carried out in a laminar flow hood.

1. Trypsinize cells as usual. Transfer 100  $\mu$ l of cell suspension to 10 ml of isotone. Count cells with the Coulter Counter.
2. Dilute cells in growth medium to 60,000 5 cells/ml. Transfer 100  $\mu$ l of cells to each well in a 96-well flat bottom plate to give 6000 cells/well.
3. Use half of plate (4 rows) for each compound and quadruplicate wells for each compound concentration, a set of 4 wells for medium control and 4 wells for DMSO 10 control.
4. Gently shake plates to allow for uniform attachment of the cells.
5. Incubate the plates at 37°C in a 10% CO<sub>2</sub> incubator.

15

Day 1: Addition of Compound: same as for Assay 1.

Day 5: Development of Assay: same as for Assay 1.

20 Assay 3: 3T3/PDGF- $\beta$ R/PDGF-BB(T) Cell Growth SRB Assay  
Cell line and growth medium:

3T3/PDGF- $\beta$ R/PDGF-BB(T) (NIH 3T3 clone 7 cells expressing PDGF  $\beta$ -receptor and PDGF-BB, from tumors resected from athymic mice)

25 2% calf serum/DMEM + 2 mM glutamine

Protocol:

Day 0: Cell Plating:

This part of assay is carried out in a laminar flow hood.

30

1. Trypsinize cells as usual. Transfer 200  $\mu$ l of cell suspension to 10 ml of isotone. Count cells on the Coulter Counter.
2. Dilute cells in growth medium to 60,000 35 cells/ml. Transfer 100  $\mu$ l of cells to each well in a 96-well flat bottom plate to give 6000 cells/well.

3. Allow half of plate (4 rows) for each compound and quadruplicate wells for each compound concentration, a set of 4 wells for medium control and 4 wells for DMSO control.

5 4. Gently shake plates to allow for uniform attachment of the cells to the plate.

5. Incubate the plates at 37°C in a 10% CO<sub>2</sub> incubator.

10 Day 1: Addition of Compound: same as for Assay 1.

Day 5: Development of Assay: same as for Assay 1.

Assay 4: Human Smooth Muscle Cells (SMC) Growth SRB Assay

15 Materials and Reagents same as for Assay 1:

Cell line and growth medium:

Human Aortic Smooth Muscle cells (Clonetics)

20 Clonetics's Bullet Kit: Smooth Muscle Basal Medium (SMBM) which is modified MCDB 131 containing fetal bovine serum (5%), hFGF (2ng/ml), hEGF (0.1 ng/ml), insulin (5.0 ug/ml), gentamicin (50ug/ml) and amphotericin B (50 ng/ml)

25 Protocol:

Day 0: Cell plating:

This part of assay is carried out in a laminar flow hood.

30 1. Trypsinize cells as usual. Transfer 200 µl of cell suspension to 10 ml of isotone. Count cells on the Coulter Counter.

2. Dilute cells in growth medium to 20,000 cells/ml. Transfer 100 µl of cells to each well in a 96-  
35 well flat bottom plate to give 2000 cells/well.

3. Allow half of plate (4 rows) for each compound and quadruplicate wells for each compound concentration,

a set of 4 wells for medium control and 4 wells for DMSO control.

4. Gently shake plates to allow for uniform attachment of the cells to the plate.
5. Incubate the plates at 37°C in a 10% CO<sub>2</sub> incubator.

Day 1: Addition of Compound: same as for Assay 1.

Day 5: Development of Assay: same as for Assay 1.

#### 6.2.3. 3T3 Cell Growth Assay

##### Assay 1: PDGF-Induced BrdU Incorporation Assay

##### Materials and Reagents:

- 15 (1) PDGF: human PDGF B/B; 1276-956, Boehringer Mannheim, Germany
- (2) BrdU Labeling Reagent: 10 mM, in PBS (pH 7.4), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- 20 (3) FixDenat: fixation solution (ready to use), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (4) Anti-BrdU-POD: mouse monoclonal antibody conjugated with peroxidase, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- 25 (5) TMB Substrate Solution: tetramethylbenzidine (TMB), ready to use, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (6) PBS Washing Solution : 1X PBS, pH 7.4, made in house.
- 30 (7) Albumin, Bovine (BSA): fraction V powder; A-8551, Sigma Chemical Co., USA.

##### Protocol

- 35 (1) 3T3 engineered cell line: 3T3/EGFRc7.



- (2) Cells are seeded at 8000 cells/well in DMEM, 10% CS, 2mM Gln in a 96 well plate. Cells are incubated overnight at 37°C in 5% CO<sub>2</sub>.
- 5 (3) After 24 hours, the cells are washed with PBS, and then are serum starved in serum free medium (0%CS DMEM with 0.1% BSA) for 24 hours.
- 10 (4) On day 3, ligand (PDGF=3.8 nM, prepared in DMEM with 0.1% BSA) and test compounds are added to the cells simultaneously. The negative control wells receive serum free DMEM with 0.1% BSA only; the positive control cells receive the ligand (PDGF) but no test compound. Test compounds are prepared in serum free DMEM with ligand in a 96 well plate, and serially
- 15 diluted for 7 test concentrations.
- (5) After 20 hours of ligand activation, diluted BrdU labeling reagent (1:100 in DMEM, 0.1% BSA) is added and the cells are incubated with BrdU (final concentration=10 µM) for 1.5
- 20 hours.
- (6) After incubation with labeling reagent, the medium is removed by decanting and tapping the inverted plate on a paper towel. FixDenat solution is added (50 µl/well) and the plates
- 25 are incubated at room temperature for 45 minutes on a plate shaker.
- (7) The FixDenat solution is thoroughly removed by decanting and tapping the inverted plate on a paper towel. Milk is added (5% dehydrated milk in PBS, 200 µl/well) as a blocking solution and
- 30 the plate is incubated for 30 minutes at room temperature on a plate shaker.
- (8) The blocking solution is removed by decanting and the wells are washed once with PBS. Anti-BrdU-POD solution (1:100 dilution in PBS, 1% BSA) is added (100 µl/well) and the plate is
- 35

incubated for 90 minutes at room temperature on a plate shaker.

- 5 (9) The antibody conjugate is thoroughly removed by decanting and rinsing the wells 5 times with PBS, and the plate is dried by inverting and tapping on a paper towel.
- 10 (10) TMB substrate solution is added (100  $\mu$ l/well) and incubated for 20 minutes at room temperature on a plate shaker until color development is sufficient for photometric detection.
- 15 (11) The absorbance of the samples are measured at 410 nm (in "dual wavelength" mode with a filter reading at 490 nm, as a reference wavelength) on a Dynatech ELISA plate reader.

#### Assay 2: EGF-Induced BrdU Incorporation Assay

##### Materials and Reagents

- 20 (1) EGF: mouse EGF, 201; Toyobo, Co., Ltd. Japan
- (2) BrdU Labeling Reagent: 10 mM, in PBS (pH7.4), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- 25 (3) FixDenat: fixation solution (ready to use), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (4) Anti-BrdU-POD: mouse monoclonal antibody conjugated with peroxidase, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- 30 (5) TMB Substrate Solution: tetramethylbenzidine (TMB), ready to use, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (6) PBS Washing Solution : 1X PBS, pH 7.4, made in house.
- 35 (7) Albumin, Bovine (BSA): fraction V powder; A-8551, Sigma Chemical Co., USA.

Protocol

- (1) 3T3 engineered cell line: 3T3/EGFRc7
- (2) Cells are seeded at 8000 cells/well in 10% CS, 2mM Gln in DMEM, in a 96 well plate.  
5 Cells are incubated overnight at 37°C in 5% CO<sub>2</sub>.
- (3) After 24 hours, the cells are washed with PBS, and then are serum starved in serum free medium (0%CS DMEM with 0.1% BSA) for 24 hours.
- (4) On day 3, ligand (EGF=2 nM, prepared in  
10 DMEM with 0.1% BSA) and test compounds are added to the cells simultaneously. The negative control wells receive serum free DMEM with 0.1% BSA only; the positive control cells receive the ligand (EGF) but no test compound.  
15 Test compounds are prepared in serum free DMEM with ligand in a 96 well plate, and serially diluted for 7 test concentrations.
- 5) After 20 hours of ligand activation,  
20 diluted BrdU labeling reagent (1:100 in DMEM, 0.1% BSA) is added and the cells are incubated with BrdU (final concentration=10 µM) for 1.5 hours.
- 6) After incubation with labeling reagent, the  
25 medium is removed by decanting and tapping the inverted plate on a paper towel. FixDenat solution is added (50 µl/well) and the plates are incubated at room temperature for 45 minutes on a plate shaker.
- (7) The FixDenat solution is thoroughly removed  
30 by decanting and tapping the inverted plate on a paper towel. Milk is added (5% dehydrated milk in PBS, 200 µl/well) as a blocking solution and the plate is incubated for 30 minutes at room temperature on a plate shaker.
- 35 (8) The blocking solution is removed by decanting and the wells are washed once with PBS. Anti-BrdU-POD solution (1:100 dilution in PBS, 1%

- BSA) is added (100  $\mu$ l/well) and the plate is incubated for 90 minutes at room temperature on a plate shaker.
- 5 (9) The antibody conjugate is thoroughly removed by decanting and rinsing the wells 5 times with PBS, and the plate is dried by inverting and tapping on a paper towel.
- 10 (10) TMB substrate solution is added (100  $\mu$ l/well) and incubated for 20 minutes at room temperature on a plate shaker until color development is sufficient for photometric detection.
- 15 (11) The absorbance of the samples are measured at 410 nm (in "dual wavelength" mode with a filter reading at 490 nm, as a reference wavelength) on a Dynatech ELISA plate reader.

**Assay 3: EGF-Induced Her2 -Driven BrdU Incorporation**  
**Materials and Reagents:**

- 20 (1) EGF: mouse EGF, 201; Toyobo, Co., Ltd. Japan
- (2) BrdU Labeling Reagent: 10 mM, in PBS (pH7.4), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- 25 (3) FixDenat: fixation solution (ready to use), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (4) Anti-BrdU-POD: mouse monoclonal antibody conjugated with peroxidase, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- 30 (5) TMB Substrate Solution: tetramethylbenzidine (TMB), ready to use, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (6) PBS Washing Solution : 1X PBS, pH 7.4, made in house.
- 35 (7) Albumin, Bovine (BSA): fraction V powder; A-8551, Sigma Chemical Co., USA.

Protocol:

- (1) 3T3 engineered cell line:  
3T3/EGFr/Her2/EGFr (EGFr with a Her2 kinase domain)
- 5 (2) Cells are seeded at 8000 cells/well in DMEM, 10% CS, 2mM Gln in a 96- well plate. Cells are incubated overnight at 37° in 5% CO<sub>2</sub>.
- (3) After 24 hours, the cells are washed with PBS, and then are serum starved in serum free  
10 medium (0%CS DMEM with 0.1% BSA) for 24 hours.
- (4) On day 3, ligand (EGF=2 nM, prepared in DMEM with 0.1% BSA) and test compounds are added to the cells simultaneously. The  
15 negative control wells receive serum free DMEM with 0.1% BSA only; the positive control cells receive the ligand (EGF) but no test compound. Test compounds are prepared in serum free DMEM with ligand in a 96 well plate, and serially diluted for 7 test concentrations.
- 20 5) After 20 hours of ligand activation, diluted BrdU labeling reagent (1:100 in DMEM, 0.1% BSA) is added and the cells are incubated with BrdU (final concentration=10 µM) for 1.5 hours.
- 25 (6) After incubation with labeling reagent, the medium is removed by decanting and tapping the inverted plate on a paper towel. FixDenat solution is added (50 µl/well) and the plates are incubated at room temperature for 45  
30 minutes on a plate shaker.
- (7) The FixDenat solution is thoroughly removed by decanting and tapping the inverted plate on a paper towel. Milk is added (5% dehydrated milk in PBS, 200 µl/well) as a blocking solution and  
35 the plate is incubated for 30 minutes at room temperature on a plate shaker.

- 5 (8) The blocking solution is removed by decanting and the wells are washed once with PBS. Anti-BrdU-POD solution (1:100 dilution in PBS, 1% BSA) is added (100  $\mu$ l/well) and the plate is incubated for 90 minutes at room temperature on a plate shaker.
- 10 (9) The antibody conjugate is thoroughly removed by decanting and rinsing the wells 5 times with PBS, and the plate is dried by inverting and tapping on a paper towel.
- 15 (10) TMB substrate solution is added (100  $\mu$ l/well) and incubated for 20 minutes at room temperature on a plate shaker until color development is sufficient for photometric detection.
- 20 (11) The absorbance of the samples are measured at 410 nm (in "dual wavelength" mode with a filter reading at 490 nm, as a reference wavelength) on a Dynatech ELISA plate reader.

**Assay 4: IGF1-Induced BrdU Incorporation Assay**  
**Materials and Reagents:**

- 25 (1) IGF1 Ligand: human, recombinant; G511, Promega Corp, USA.
- (2) BrdU Labeling Reagent: 10 mM, in PBS (pH7.4), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- 30 (3) FixDenat: fixation solution (ready to use), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- (4) Anti-BrdU-POD: mouse monoclonal antibody conjugated with peroxidase, Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- 35 (5) TMB Substrate Solution: tetramethylbenzidine (TMB), ready to use, Cat. No. 1 647 229, Boehringer Mannheim, Germany.

- (6) PBS Washing Solution : 1X PBS, pH 7.4, made in house.
- (7) Albumin, Bovine (BSA): fraction V powder; A-8551, Sigma Chemical Co., USA.

5

Protocol:

- (1) 3T3 engineered cell line: 3T3/IGF1r.
- (2) Cells are seeded at 8000 cells/well in DMEM, 10% CS, 2mM Gln in a 96- well plate.
- 10 Cells are incubated overnight at 37°C in 5% CO<sub>2</sub>.
- (3) After 24 hours, the cells are washed with PBS, and then are serum starved in serum free medium (0%CS DMEM with 0.1% BSA) for 24 hours.
- (4) On day 3, ligand (IGF1=3.3 nM, prepared in DMEM with 0.1% BSA) and test compounds are added to the cells simultaneously. The negative control wells receive serum free DMEM with 0.1% BSA only; the positive control cells receive the ligand (IGF1) but no test compound.
- 15 Test compounds are prepared in serum free DMEM with ligand in a 96.well plate, and serially diluted for 7 test concentrations.
- 20
- 5) After 16 hours of ligand activation, diluted BrdU labeling reagent (1:100 in DMEM, 0.1% BSA) is added and the cells are incubated with BrdU (final concentration=10 µM) for 1.5 hours.
- 25
- (6) After incubation with labeling reagent, the medium is removed by decanting and tapping the inverted plate on a paper towel. FixDenat solution is added (50 µl/well) and the plates are incubated at room temperature for 45 minutes on a plate shaker.
- 30
- (7) The FixDenat solution is thoroughly removed by decanting and tapping the inverted plate on a paper towel. Milk is added (5% dehydrated milk in PBS, 200 µl/well) as a blocking solution and
- 35

- the plate is incubated for 30 minutes at room temperature on a plate shaker.
- 5 (8) The blocking solution is removed by decanting and the wells are washed once with PBS. Anti-BrdU-POD solution (1:100 dilution in PBS, 1% BSA) is added (100  $\mu$ l/well) and the plate is incubated for 90 minutes at room temperature on a plate shaker.
- 10 (9) The antibody conjugate is thoroughly removed by decanting and rinsing the wells 5 times with PBS, and the plate is dried by inverting and tapping on a paper towel.
- 15 (10) TMB substrate solution is added (100  $\mu$ l/well) and incubated for 20 minutes at room temperature on a plate shaker until color development is sufficient for photometric detection.
- 20 (11) The absorbance of the samples are measured at 410 nm (in "dual wavelength" mode with a filter reading at 490 nm, as a reference wavelength) on a Dynatech ELISA plate reader.

**Assay 5: Insulin-Induced BrdU Incorporation Assay**  
**Materials and Reagents:**

- 25 (1) Insulin: crystalline, bovine, Zinc; 13007, Gibco BRL, USA.
- (2) BrdU Labeling Reagent: 10 mM, in PBS (pH7.4), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- 30 (3) FixDenat: fixation solution (ready to use), Cat. No. 1 647 229, Boehringer Mannheim, Germany.
- 35 (4) Anti-BrdU-POD: mouse monoclonal antibody conjugated with peroxidase, Cat. No. 1 647 229, Boehringer Mannheim, Germany.



## (5) TMB Substrate Solution:

tetramethylbenzidine (TMB), ready to use, Cat. No. 1 647 229, Boehringer Mannheim, Germany.

## (6) PBS Washing Solution : 1X PBS, pH 7.4, made in house.

## (7) Albumin, Bovine (BSA): fraction V powder; A-8551, Sigma Chemical Co., USA.

Protocol:

- 10 (1) 3T3 engineered cell line: H25
- (2) Cells are seeded at 8000 cells/well in DMEM, 10% CS, 2mM Gln in a 96 well plate. Cells are incubated overnight at 37°C in 5% CO<sub>2</sub>.
- 15 (3) After 24 hours, the cells are washed with PBS, and then are serum starved in serum free medium (0%CS DMEM with 0.1% BSA) for 24 hours.
- (4) On day 3, ligand (Insulin=10 nM, prepared in DMEM with 0.1% BSA) and test compounds are added to the cells simultaneously. The
- 20 negative control wells receive serum free DMEM with 0.1% BSA only; the positive control cells receive the ligand (Insulin) but no test compound. Test compounds are prepared in serum free DMEM with ligand in a 96 well plate, and
- 25 serially diluted for 7 test concentrations.
- (5) After 16 hours of ligand activation, diluted BrdU labeling reagent (1:100 in DMEM, 0.1% BSA) is added and the cells are incubated with BrdU (final concentration=10 µM) for 1.5
- 30 hours.
- (6) After incubation with labeling reagent, the medium is removed by decanting and tapping the inverted plate on a paper towel. FixDenat solution is added (50 µl/well) and the plates
- 35 are incubated at room temperature for 45 minutes on a plate shaker.

- 5 (7) The FixDenat solution is thoroughly removed by decanting and tapping the inverted plate on a paper towel. Milk is added (5% dehydrated milk in PBS, 200  $\mu$ l/well) as a blocking solution and the plate is incubated for 30 minutes at room temperature on a plate shaker.
- 10 (8) The blocking solution is removed by decanting and the wells are washed once with PBS. Anti-BrdU-POD solution (1:100 dilution in PBS, 1% BSA) is added (100  $\mu$ l/well) and the plate is incubated for 90 minutes at room temperature on a plate shaker.
- 15 (9) The antibody conjugate is thoroughly removed by decanting and rinsing the wells 5 times with PBS, and the plate is dried by inverting and tapping on a paper towel.
- 20 (10) TMB substrate solution is added (100  $\mu$ l/well) and incubated for 20 minutes at room temperature on a plate shaker until color development is sufficient for photometric detection.
- 25 (11) The absorbance of the samples are measured at 410 nm (in "dual wavelength" mode with a filter reading at 490 nm, as a reference wavelength) on a Dynatech ELISA plate reader.

#### 6.2.4. HUV-EC-C Assay

The following protocol may also be used to measure a compound's activity:

##### 30 DAY 0

1. Wash and trypsinize HUV-EC-C cells (human umbilical vein endothelial cells, (American Type Culture Collection; catalogue no. 1730 CRL). Wash with Dulbecco's phosphate-buffered saline (D-PBS; obtained 35 from Gibco BRL; catalogue no. 14190-029) 2 times at about 1 ml/10 cm<sup>2</sup> of tissue culture flask. Trypsinize with 0.05% trypsin-EDTA in non-enzymatic cell dissociation

solution (Sigma Chemical Company; catalogue no. C-1544). The 0.05% trypsin was made by diluting 0.25% trypsin/1 mM EDTA (Gibco; catalogue no. 25200-049) in the cell dissociation solution. Trypsinize with about 1 ml/25-30 cm<sup>2</sup> of tissue culture flask for about 5 minutes at 37°C. After cells have detached from the flask, add an equal volume of assay medium and transfer to a 50 ml sterile centrifuge tube (Fisher Scientific; catalogue no. 05-539-6).

10 2. Wash the cells with about 35 ml assay medium in the 50 ml sterile centrifuge tube by adding the assay medium, centrifuge for 10 minutes at approximately 200xg, aspirate the supernatant, and resuspend with 35 ml D-PBS. Repeat the wash two more times with D-PBS, resuspend the  
15 cells in about 1 ml assay medium/15 cm<sup>2</sup> of tissue culture flask. Assay medium consists of F12K medium (Gibco BRL; catalogue no. 21127-014) + 0.5% heat-inactivated fetal bovine serum. Count the cells with a Coulter Counter®v Coulter Electronics, Inc.) and add assay medium to the  
20 cells to obtain a concentration of 0.8-1.0x10<sup>5</sup> cells/ml.

3. Add cells to 96-well flat-bottom plates at 100 µl/well or 0.8-1.0x10<sup>4</sup> cells/well; incubate -24h at 37°C, 5% CO<sub>2</sub>.

#### DAY 1

25 1. Make up two-fold drug titrations in separate 96-well plates, generally 50 µM on down to 0 µM. Use the same assay medium as mentioned in day 0, step 2 above. Titrations are made by adding 90 µl/well of drug at 200 µM (4X the final well concentration) to the top well of a  
30 particular plate column. Since the stock drug concentration is usually 20 mM in DMSO, the 200 µM drug concentration contains 2% DMSO.

Therefore, diluent made up to 2% DMSO in assay medium (F12K + 0.5% fetal bovine serum) is used as  
35 diluent for the drug titrations in order to dilute the drug but keep the DMSO concentration constant. Add this diluent to the remaining wells in the column at 60

$\mu$ l/well. Take 60  $\mu$ l from the 120  $\mu$ l of 200  $\mu$ M drug dilution in the top well of the column and mix with the 60  $\mu$ l in the second well of the column. Take 60  $\mu$ l from this well and mix with the 60  $\mu$ l in the third well of the column, and so on until two-fold titrations are completed. When the next-to-the-last well is mixed, take 60  $\mu$ l of the 120  $\mu$ l in this well and discard it. Leave the last well with 60  $\mu$ l of DMSO/media diluent as a non-drug-containing control. Make 9 columns of titrated drug, enough for triplicate wells each for 1) VEGF (obtained from Pepro Tech Inc., catalogue no. 100-200; 2) endothelial cell growth factor (ECGF) (also known as acidic fibroblast growth factor, or aFGF) (obtained from Boehringer Mannheim Biochemica, catalogue no. 1439 600), and assay media control. ECGF comes as a preparation with sodium heparin.

2. Transfer 50  $\mu$ l/well of the drug dilutions to the 96-well assay plates containing the  $0.8-1.0 \times 10^4$  cells/100  $\mu$ l/well of the HUV-EC-C cells from day 0 and incubate ~2 h at 37°C, 5% CO<sub>2</sub>.

3. In triplicate, add 50  $\mu$ l/well of 80  $\mu$ g/ml VEGF, 20 ng/ml ECGF, or media control to each drug condition. As with the drugs, the growth factor concentrations are 4X the desired final concentration. Use the assay media from day 0 step 2 to make the concentrations of growth factors. Incubate approximately 24 hours at 37°C, 5% CO<sub>2</sub>. Each well will have 50  $\mu$ l drug dilution, 50  $\mu$ l growth factor or media, and 100  $\mu$ l cells, = 200  $\mu$ l/well total. Thus the 4X concentrations of drugs and growth factors become 1X once everything has been added to the wells.

#### DAY 2

1. Add <sup>3</sup>H-thymidine (Amersham; catalogue no. TRK-686) at 1  $\mu$ Ci/well (10  $\mu$ l/well of 100  $\mu$ Ci/ml solution made up in RPMI media + 10% heat-inactivated fetal bovine serum) and incubate ~24 h at 37°C, 5% CO<sub>2</sub>. Note: <sup>3</sup>H-thymidine is made up in RPMI media because all of the other applications for which we use the <sup>3</sup>H-thymidine

involve experiments done in RPMI. The media difference at this step is probably not significant. RPMI was obtained from Gibco BRL, catalogue no. 11875-051.

DAY 3

- 5 1. Freeze plates overnight at  $-20^{\circ}\text{C}$ .

DAY 4

1. Thaw plates and harvest with a 96-well plate harvester (Tomtec Harvester 96<sup>(R)</sup>) onto filter mats (Wallac; catalogue no. 1205-401); read counts on a Wallac 10 Betaplate<sup>(TM)</sup> liquid scintillation counter.

6.2.5. PDGF-R Cellular Assay

The PDGF cellular kinase assay was carried out as follows: cells are lysed in 0.2 M Hepes, 0.15 M NaCl, 10% V/V glycerol, 0.04% Triton X-100, 5 mM EDTA, 5 mM sodium vanadate and 2 mM Na<sup>+</sup> pyrophosphate; cell lysates are then added to an ELISA plate coated with an anti-PDGF receptor antibody (Genzyme); ELISA plates are coated at 0.5  $\mu\text{g}$  of antibody/well in 150  $\mu\text{l}$  of PBS for 18 hours at 4 $^{\circ}\text{C}$  prior to the addition of the lysate; the lysate is incubated in the coated plates for 1 hour and then washed four times in TBST (35 mM Tris-HCl pH 7.0, 0.15 M NaCl, 0.1% Triton X100); anti-phosphotyrosine antibody (100  $\mu\text{l}$  in PBS) is added and the mixture is incubated for 30 minutes at room temperature; the wells were then washed four times in TBST, a secondary antibody conjugated to POD (TAGO) is added to each well, and the treated well are incubated for 30 minutes at room temperature; the wells are then washed four times in TBST, ABTS/H<sub>2</sub>O<sub>2</sub> solution is added to each well and the wells are incubated for two minutes; absorbance is then measured at 410 nm.

### 6.2.6. Experimental Results Of Cell Growth Assay

Results for various compounds obtained from the above-described assays are set forth in the  
5 Tables that follow:

TABLE 2

Mitogenesis in Endothelial Cells  
[3H]Thymidine Incorporation

	COMPOUND	HUV-EC	Assay
		VEGF ( $\mu$ M)	a-FGF ( $\mu$ M)
10	SU4312	1.1	153.8
	SU4314	0.2	6.0
	SU4793	6.6	3.4
15	SU4794	4.8	35.7
	SU4796	30.7	35.8
	SU4798	43.2	
	SU4799	19.9	
	SU4932	2.5	45.2
20	SU4942	1.6	4.6
	SU4944	14.8	
	SU4949	3.4	3.7
	SU4952	25.6	19.3
25	SU4956	8.0	13.0
	SU4967	34.3	16.3
	SU4972	1.0	1.4
	SU4979	4.4	4.9
	SU4981	0.6	
30	SU4982	46.1	27.3
	SU4984	0.8	25.8
	SU5201	2.5	2.3
	SU5204	2.3	0.7
35	SU5205	5.1	11.8
	SU5208	2.9	130
	SU5217	9.6	10.5

5

10

15

20

25

30

35

COMPOUND	HUV-EC VEGF ( $\mu$ M)	Assay a-FGF ( $\mu$ M)
SU5218	2.4	2.7
SU5401	2.2	
SU5402	<0.8	2.0
SU5404	<0.8	31.1
SU5405	0.9	0.6
SU5406	<0.8	
SU5407	39.8	35.5
SU5408	<0.8	22.7
SU5409	26.0	
SU5416	<0.8	
SU5418	13.6	40
SU5419	0.7	
SU5421	11.4	
SU5424	2.5	
SU5427	5.7	
SU5429	27.6	
SU5432	0.16	0.14
SU5438	39.8	33.0
SU5451	1.2	30.0
SU5454	3.8	3.4
SU5455	20	20
SU5461	<0.07	<0.07
SU5462	0.5	0.8
SU5463	0.14	7.9
SU5464	3.8	12.9
SU5466	1.3	3.2
SU5468	0.54	8.7
SU5472	2.0	5.0
SU5473	1.2	14.1
SU5477	0.05	37.8

COMPOUND	HUV-EC Assay	
	VEGF ( $\mu\text{M}$ )	a-FGF ( $\mu\text{M}$ )
SU5480	1.2	3.8

5

**TABLE 3**  
Mitogenesis in 3T3/EGFR Cells  
BrdU Incorporation

10

15

20

25

30

35

COMPOUND	PDGFR PDGF Ligand IC50 ( $\mu\text{M}$ )	FGFR FGF Ligand IC50 ( $\mu\text{M}$ )	EGFR EGF Ligand IC50 ( $\mu\text{M}$ )
SU4312	75		
SU4313	6	5.5	5.5
SU4314	2.5		
SU4967	9	4.9	60
SU4981	3	10	20
SU5402	50	40	
SU5404	3	25	
SU5406	5.2		
SU5407	7.5	70	100
SU5416	2.8	70	
SU5451	30	16	
SU5463			23
SU5464	70	60	95
SU5465	40	25	50
SU5466	18	15	17
SU5468	8		
SU5469	4	15	28
SU5473	4	50	54
SU5475	6.5	9	48



TABLE 4  
Cell Growth Assay on Various Cell Lines

SRB Readout

COMPOUND	3T3/E/H+ TGF- $\alpha$ (T) IC50 ( $\mu$ M)	3T3/EGFR+ TGF- $\alpha$ (T) IC50 ( $\mu$ M)	3T3/PDGFR+ PDGF(T) IC50 ( $\mu$ M)	SMC IC50 ( $\mu$ M)
SU4312	36			
SU4313	32	10.7		8.8
SU4314	78		10	
SU4984			22.2	

3T3/E/H+TGF- $\alpha$ (T): NIH 3T3 cells expressing  
EGFR/HER2 chimera and TGF- $\alpha$ , tumor-derived

3T3/EGFR+TGF- $\alpha$ (T): NIH 3T3 cells expressing EGFR and  
TGF- $\alpha$ , tumor-derived

3T3/PDGFR+PDGF(T): NIH 3T3 cells expressing PDGF- $\beta$ R  
and PDGF- $\beta\beta$ , tumor-derived

SMC: human smooth muscle cells from Clonetics

### 6.3. Measurement Of Cell Toxicity

Therapeutic compounds should be more potent in  
inhibiting receptor tyrosine kinase activity than in  
exerting a cytotoxic effect. A measure of the  
effectiveness and cell toxicity of a compound can be  
obtained by determining the therapeutic index:  $IC_{50}/LD_{50}$ .  
 $IC_{50}$ , the dose required to achieve 50% inhibition, can be  
measured using standard techniques such as those  
described herein.  $LD_{50}$ , the dosage which results in 50%  
toxicity, can also be measured by standard techniques  
(Mossman, 1983, *J. Immunol. Methods*, 65:55-63), by  
measuring the amount of LDH released (Korzeniewski and  
Callewaert, 1983, *J. Immunol. Methods* 64:313; Decker and  
Lohmann-Matthes, 1988, *J. Immunol. Methods* 115:61), or by  
measuring the lethal dose in animal models. Compounds  
with a large therapeutic index are preferred. The  
therapeutic index should be greater than 2, preferably at  
least 10, more preferably at least 50.

### 6.3. In Vivo Animal Models

#### 6.3.1. Xenograft Animal Models

The ability of human tumors to grow as xenografts in athymic mice (e.g., Balb/c, nu/nu) provides a useful in vivo model for studying the biological response to therapies for human tumors. Since the first successful xenotransplantation of human tumors into athymic mice, (Rygaard and Povlsen, 1969, *Acta Pathol. Microbial. Scand.* 77:758-760), many different human tumor cell lines (e.g., mammary, lung, genitourinary, gastrointestinal, head and neck, glioblastoma, bone, and malignant melanomas) have been transplanted and successfully grown in nude mice. Human mammary tumor cell lines, including MCF-7, ZR75-1, and MDA-MB-231, have been established as subcutaneous xenografts in nude mice (Warri et al., 1991, *Int. J. Cancer* 49:616-623; Ozzello and Sordat, 1980, *Eur. J. Cancer* 16:553-559; Osborne et al., 1985, *Cancer Res.* 45:584-590; Seibert et al., 1983, *Cancer Res.* 43:2223-2239).

#### Assay 1: HER2/Xenograft Animal Model

To study the effect of anti-tumor drug candidates on HER2 expressing tumors, the tumor cells should be able to grow in the absence of supplemental estrogen. Many mammary cell lines are dependent on estrogen for in vivo growth in nude mice (Osborne et al., *supra*), however, exogenous estrogen suppresses HER2 expression in nude mice (Warri et al., *supra*, Dati et al., 1990, *Oncogene* 5:1001-1006). For example, in the presence of estrogen, MCF-7, ZR-75-1, and T47D cells grow well in vivo, but express very low levels of HER2 (Warri et al., *supra*, Dati et al., *supra*).

The following type of xenograft protocol can be used:

- 1) implant tumor cells (subcutaneously) into the hindflank of five- to six-week-old female Balb/c nu/nu athymic mice;

- 2) administer the anti-tumor compound;
- 3) measure tumor growth by measuring tumor volume.

The tumors can also be analyzed for the presence of a receptor, such as HER2, EGF or PDGF, by Western and immunohistochemical analyses. Using techniques known in the art, one skilled in the art can vary the above procedures, for example through the use of different treatment regimes.

**Assay 2: FLK-1/Xenograft Model.**

10 The ability of the compounds of the present invention to inhibit ovarian, melanoma, prostate, lung and mammary tumor cell lines established as SC xenografts was examined. These studies were conducted using doses ranging from 1 to 75 mg/kg/day.

15 **Materials And Methods.** The tumor cells were implanted subcutaneously into the indicated strains of mice. Treatment was initiated on day 1 post implantation unless otherwise indicated (e.g. treatment of the SCID mouse related to the A375 melanoma cell line began on Day 20 9). Eight (8) to sixteen (16) mice comprised each test group.

**Specifically:**

**Animals.** Female athymic mice (BALB/c, nu/nu), BALB/c mice, Wistar rats and Fisher 344 rats were 25 obtained from Simonsen Laboratories (Gilroy, CA). Female A/I mice were obtained from Jackson Laboratory (Bar Harbor, ME). DA rats were obtained from B&K Universal, Inc. (Fremont, CA). Athymic R/Nu rats, DBA/2N mice, and BALB/c mice were obtained from Harlan Sprague Dawley 30 (Indianapolis, IN). Female C57BL/6 mice were obtained from Taconic (Germantown, NY). All animals were maintained under clean-room conditions in Micro-isolator cages with Alpha-dri bedding. They received sterile rodent chow and water *ad libitum*.

35 All procedures were conducted in accordance with the NIH Guide for the Care and Use Of Laboratory Animals.

*Subcutaneous Xenograft Model.* Cell lines were grown in appropriate medium as described (See Section 6). Cells were harvested at or near confluency with 0.05% Trypsin-EDTA and pelleted at 450 x g for 10 min. Pellets were resuspended in sterile PBS or media (without FBS) to a suitable concentration indicated in the Figure legends and the cells were implanted into the hindflank of mice. Tumor growth was measured over 3 to 6 weeks using vernier calipers tumor volumes were calculated as a product of length x width x height unless otherwise indicated. P values were calculated using the Students' t-test. Compound in 50 - 100  $\mu$ L excipient (dimethylsulfoxide, PBTE, PBTE6C:D5W, or PBTE:D5W) was delivered by IP injection at concentrations indicated in the Figure legends.

*Intracerebral Xenograft Model.* For the mouse IC model, rat C6 glioma cells were harvested and suspended in sterile PBS at a concentration of  $2.5 \times 10^7$  cells/ml and placed on ice. Cells were implanted into BALB/c, nu/nu mice in the following manner: the frontoparietal scalps of mice were shaved with animal clippers if necessary before swabbing with 70% ethanol. Animals were anesthetized with isofluorane and the needle was inserted through the skull into the left hemisphere of the brain. Cells were dispensed from Hamilton Gas-tight Syringes using 30 ga 1/2 inch needles fitted with sleeves that allowed only a 3 mm penetration. A repeater dispenser was used for accurate delivery of 4  $\mu$ L of cell suspension. Animals were monitored daily for well-being and were sacrificed when they had a weight loss of about 40% and/or showed neurological symptoms.

For the rat IC model, rats (Wistar, Sprague Dawley, Fisher 344, or athymic R/Nu; approximately 200-400 g (some 3-400g)) were anesthetized by an IP injection of 100 mg/kg Ketaset (ketamine hydrochloride; Avaco, Fort Dodge, Iowa) and 5 mg/kg Rompun (xylazine, 2% solution; Bayer, Germany). After onset of anesthesia, the scalp

was shaved and the animal was oriented in a stereotaxic apparatus (Stoelting, Wood Dale, IL). The skin at the incision site was cleaned 3 times with alternating swabs of 70% ethanol and 10% Povidone-Iodine. A median 1.0 - 5 1.5 cm incision was made in the scalp using a sterile surgical blade. The skin was detached slightly and pulled to the sides to expose the sutures on the skull surface. A dental drill (Stoelting, Wood Dale, IL) was used to make a small (1-2 mm diameter) burrhole in the 10 skull approximately 1 mm anterior and 2 mm lateral to the bregma. The cell suspension was drawn into a 50  $\mu$ L Hamilton syringe fitted with a 23 or 25g standard bevel needle. The syringe was oriented in the burrhole at the level of the arachnoidea and lowered until the tip of the 15 needle was 3 mm deep into the brain structure, where the cell suspension was slowly injected. After cells were injected, the needle was left in the burrhole for 1-2 minutes to allow for complete delivery of the cells. The skull was cleaned and the skin was closed with 2 to 3 20 sutures. Animals were observed for recovery from surgery and anesthesia. Throughout the experiment, animals were observed at least twice each day for development of symptoms associated with progression of intracerebral tumor. Animals displaying advanced symptoms (leaning, 25 loss of balance, dehydration, loss of appetite, loss of coordination, cessation of grooming activities, and/or significant weight loss) were humanely sacrificed and the organs and tissues of interest were resected.

*Intraperitoneal Model.* Cell lines were grown 30 in the appropriate media. Cells were harvested and washed in sterile PBS or medium without FBS, resuspended to a suitable concentration, and injected into the IP cavity of mice of the appropriate strain. Mice were observed daily for the occurrence of ascites formation. 35 Individual animals were sacrificed when they presented with a weight gain of 40%, or when the IP tumor burden began to cause undue stress and pain to the animal.

### 6.3.2. In Vivo VEGF Pellet Model

In the following example, the Pellet Model was used to test a compound's activity against the FLK-1 receptor and against disorders associated with the formation of blood vessels. In this model, VEGF is packaged into a time-release pellet and implanted subcutaneously on the abdomen of nude mice to induce a 'reddening' response and subsequent swelling around the pellet. Potential FLK-1 inhibitors may then be implanted in methylcellulose near the VEGF pellet to determine whether such inhibitor may be used to inhibit the "reddening" response and subsequent swelling.

**Materials And Methods.** The following materials were used:

- 1) VEGF- human recombinant lyophilized product is commercially may be obtained from Peprotech, Inc., Princeton Business Park, G2; P.O. box 275, Rocky Hill, NJ 08553.
- 2) VEGF packaged into 21 day release pellets were obtained from Innovative Research of America (Innovative Research of America, 3361 Executive Parkway, P.O. Box 2746, Toledo, Ohio 43606), using patented matrix driven delivery system. Pellets were packaged at 0.20, 0.21, or 2.1 µg VEGF/pellet. These doses approximate 10 and 100 ng/day release of VEGF.
- 3) Methylcellulose
- 4) Water (sterile)
- 5) Methanol
- 6) Appropriate drugs/inhibitors
- 7) 10 cm culture plates
- 8) parafilm

The following protocol was then followed to conduct the VEGF pellet model:

- 1) VEGF, purchased from Peprotech, was sent to Innovative Research for Custom Pellet preparation;
- 2) Methylcellulose prepared at 1.5% (w/v) in sterile water;

- 3) Drugs solubilized in methanol (usual concentration range = 10 to 20 mg/ml);
- 4) Place sterile parafilm in sterile 10 cm plates;
- 5) 150  $\mu$ l of drug in methanol added to 1.35 ml of 1.5% methylcellulose and mixed/vortexed thoroughly;
- 6) 25  $\mu$ l aliquots of homogenate placed on parafilm and dried into discs;
- 7) Mice (6-10 wk. Balb/C athymic nu/nu, female) were anesthetized via isoflurane inhalation;
- 8) VEGF pellets and methylcellulose discs were implanted subcutaneously on the abdomen; and
- 9) Mice were scored at 24 hours and 48 hours for reddening and swelling response.

The specific experimental design used in this example was:

N = 4 animals/group

Controls: VEGF pellet + drug placebo  
VEGF placebo + drug pellet

**Experimental Results.** The compounds of the present invention are expected to demonstrate activity according to this assay.

### 6.3.3. Mammary Fat Pad Model

Because of the established role played by many of the RTKs, e.g., the HER2 receptor, in breast cancer, the mammary fat pad model is particularly useful for measuring the efficacy of compounds which inhibit such RTKs. By implanting tumor cells directly into the location of interest, *in situ* models more accurately reflect the biology of tumor development than do subcutaneous models. Human mammary cell lines, including MCF-7, have been grown in the mammary fat pad of athymic mice. Shafie and Grantham, 1981, *Natl. Cancer Instit.* 67:51-56; Gottardis et al., 1988, *J. Steroid Biochem.* 30:311-314. More specifically, the following procedure can be used to measure the inhibitory effect of a compound on the HER2 receptor:

- 1) Implant, at various concentrations, MDA-MB-231 and MCF-7 cells transfected with HER-2 into the axillary mammary fat pads of female athymic mice;
- 2) Administer the compound; and
- 3) Measure the tumor growth at various time points.

The tumors can also be analyzed for the presence of a receptor such as HER2, by Western and immunohistochemical analyses. Using techniques known in the art, one skilled in the art can vary the above procedures, for example through the use of different treatment regimes.

#### 6.3.4. Tumor Invasion Model

The following tumor invasion model has been developed and may be used for the evaluation of therapeutic value and efficacy of the compounds identified to selectively inhibit KDR/FLK-1 receptor.

##### 6.3.4.1. Procedure

8 week old nude mice (female) (Simonsen Inc.) were used as experimental animals. Implantation of tumor cells was performed in a laminar flow hood. For anesthesia, Xylazine/Ketamine Cocktail (100 mg/kg ketamine and 5 mg/kg) are administered intraperitoneally. A midline incision is done to expose the abdominal cavity (approximately 1.5 cm in length) to inject  $10^7$  tumor cells in a volume of 100  $\mu$ l medium. The cells are injected either into the duodenal lobe of the pancreas or under the serosa of the colon. The peritoneum and muscles are closed with a 6-0 silk continuous suture and the skin was closed by using wound clips. Animals were observed daily.

##### 6.3.4.2. Analysis



After 2-6 weeks, depending on gross observations of the animals, the mice are sacrificed, and the local tumor metastases, to various organs (lung, liver, brain, stomach, spleen, heart, muscle) are excised and analyzed (measurements of tumor size, grade of invasion, immunochemistry, and in situ hybridization).

#### 6.3.5. RESULTS

Results for various compounds  
10 obtained from the above-described in vivo assays are set forth at Table 5, below:

15

20

25

30

35

TABLE 5  
In Vivo Data

5	<div>COMPOUND</div> <div>Eph4-VEGF %inhibition @ mg/kg</div>
	<div>SU4312</div> <div>56% @ 75</div> <div>-----</div> <div>50% @ 75</div> <div>63% @ 50</div>
10	<div>SU4932</div> <div>42% @ 75</div> <div>-----</div> <div>--</div> <div>42% @ 50/50</div>
	<div>SU4942</div> <div>46% @ 50</div> <div>47% @ 25</div>
15	<div>SU5416</div> <div>50% @ 25</div> <div>-----</div> <div>--</div> <div>57% @ 37.5/37.5</div>
	<div>SU5424</div> <div>45% @ 50</div> <div>-----</div> <div>--</div> <div>65% @ 50</div>
20	<div>SU5427</div> <div>47% @ 50</div> <div>-----</div> <div>--</div> <div>65% @ 50</div>

The present invention is not to be limited in scope by the exemplified embodiments which are intended as illustrations of single aspects of the invention, and any clones, DNA or amino acid sequences which are functionally equivalent are within the scope of the invention. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

All references cited herein are hereby incorporated by reference in their entirety.


CLAIMS

What Is Claimed:

1. A compound of the formula

5

10



and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

15  $R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$R_3'$ ,  $R_5'$ , and  $R_6'$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$n$  is 0-3;

$X$  is Br, Cl, F or I;

$R$  is H, alkyl or aryl; and

30  $R'$  is H, alkyl or aryl.

2. The compound of claim 1 wherein:


$R_3'$  is selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where R is H, alkyl or aryl; and

$R_5'$  is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where R is H, alkyl or aryl.

5

3. A compound of the formula:

10



15

and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

20  $R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

25  $R_2'$ ,  $R_3'$ ,  $R_4'$ , and  $R_5'$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

30  $n$  is 0-3;

$R$  is H, alkyl or aryl;

$R'$  is H, alkyl or aryl; and

$R_a$  and  $R_b$  are each independently selected from the group consisting of H, alkyl and  $C(O)R$ , or  $NRaRb$  taken together are a heterocyclic ring of from 3 to 8 atoms optionally substituted at one or more positions with hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl,

alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>3</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R or CONRR';

with the provisos that:

5 (i) when Ra and Rb are both H, methyl, or ethyl, at least one of R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>3</sub>' or R<sub>5</sub>' must be other than hydrogen; and

(ii) when R<sub>1</sub> is methyl and one of Ra and Rb is methyl and the other is isooctyl, at least one of R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, 10 R<sub>3</sub>' or R<sub>5</sub>' must be other than hydrogen.

4. The compound of claim 3 wherein:


R<sub>3</sub>' is selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert- 15 butyl, halogen, aryl and OR, where R is H, alkyl or aryl; and

R<sub>5</sub>' is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where R is 20 H, alkyl or aryl.

5. A pharmaceutical composition comprising a compound of the formula:

25

30



or pharmaceutically acceptable salts thereof, in combination with a pharmaceutically acceptable carrier or 35 excipient, wherein:

R<sub>1</sub> is H or alkyl;

R<sub>2</sub> is O or S;

R<sub>1</sub> is hydrogen;

R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>7</sub> are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>3</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';

R<sub>2</sub>', R<sub>3</sub>', R<sub>5</sub>', and R<sub>6</sub>' are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>3</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';

R<sub>a</sub> and R<sub>b</sub> are each independently selected from the group consisting of H, alkyl and C(O)R, or NR<sub>a</sub>R<sub>b</sub> taken together are a heterocyclic ring of from 3 to 8 atoms optionally substituted at one or more positions with hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>3</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R or CONRR';

n is 0-3;

R is H, alkyl or aryl; and

R' is H, alkyl or aryl;


with the proviso that when the pharmaceutically acceptable carrier or excipient is selected from the group consisting of anise oil, carboxymethylcellulose and peanut oil, the compound is not 3-(4-dimethylaminobenzylidene)-2-indolinone.

6. The pharmaceutical composition of claim 5 wherein:

R<sub>1</sub>' is selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where R is H, alkyl or aryl; and

R<sub>4</sub>' is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where R is H, alkyl or aryl.

7. A compound of the formula:



and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

15  $R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

20 A is a five membered heteroaryl ring selected from the group consisting of thiophene, pyrrole, pyrazole, imidazole, 1,2,3-triazole, 1,2,4-triazole, oxazole, isoxazole, thiazole, isothiazole, 2-sulfonylfuran, 4-alkylfuran, 1,2,3-oxadiazole, 1,2,4-oxadiazole, 1,2,5-oxadiazole, 1,3,4-oxadiazole, 1,2,3,4-oxatriazole, 1,2,3,5-oxatriazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,3,4-thiatriazole, 1,2,3,5-thiatriazole, and tetrazole, optionally substituted at one or more  
25 positions with alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$  or  $CONRR'$ ;

n is 0-3;


35 R is H, alkyl or aryl; and

R' is H, alkyl or aryl;

with the proviso that the following compounds are excluded:

- 3-(pyrrol-2-ylmethylene)-2-indolinone;  
 3-(5-chloro-3,4-dimethylpyrrol-2-ylmethylene)-2-indolinone;  
 3-(3,5-dimethyl-4-ethylpyrrol-2-yl)-2-indolinone;  
 3-(3,5-dimethyl-4-ethoxycarbonylpyrrol-2-yl)-2-indolinone;  
 Benzoic acid, 2-[[[1-ethyl-2,3-dihydro-2-oxo-3-(1H-pyrrol-2-ylmethylene)-1H-indol-5-yl]oxy)methyl]-;  
 3-[(1-methyl-5-nitro-imidazol-2-yl)methylene]-2-indolinone;  
 1-methyl-3-[(1-methyl-5-nitro-imidazol-2-yl)methylene]-2-indolinone;  
 1-ethyl-3-[(1-methyl-5-nitro-imidazol-2-yl)methylene]-2-indolinone;  
 3-(thien-2-ylmethylene)-2-indolinone;  
 1H-Indole-7-acetic acid, 3-[(2-butyl-1H-imidazol-4-yl)methylene]-2,3-dihydro-2-oxo-, ethyl ester;  
 1H-Indole-7-acetic acid, 3-[[2-butyl-1-[(1,1-dimethylethoxy)carbonyl]-1H-imidazol-4-yl)methylene]-2,3-dihydro-2-oxo-, ethyl ester.

25 8. A pharmaceutical composition comprising a compound of the formula:





or pharmaceutically acceptable salts thereof, in combination with a pharmaceutically acceptable carrier or excipient, wherein:

- $R_1$  is H or alkyl;
- 5  $R_2$  is O or S;
- $R_3$  is hydrogen;
- $R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, 10 CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';
- A is a five membered heteroaryl ring selected from thiophene, pyrrole, pyrazole, imidazole, 1,2,3-triazole, 1,2,4-triazole, oxazole, isoxazole, thiazole, 15 isothiazole, 2-sulfonylfuran, 4-alkylfuran, 1,2,3-oxadiazole, 1,2,4-oxadiazole, 1,2,5-oxadiazole, 1,3,4-oxadiazole, 1,2,3,4-oxatriazole, 1,2,3,5-oxatriazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,3,4-thiatriazole, 1,2,3,5- 20 thiatriazole, and tetrazole, optionally substituted at one or more positions with alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH, CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R or CONRR';
- 25  $n$  is 0-3;
- $R$  is H, alkyl or aryl; and
- $R'$  is H, alkyl or aryl;

with the proviso that the following compounds are excluded:

- 30 3-(pyrrol-2-ylmethylene)-2-indolinone;
- 3-(5-chloro-3,4-dimethylpyrrol-2-ylmethylene)-2-indolinone;
- 3-(3,5-dimethyl-4-ethylpyrrol-2-yl)-2-indolinone;
- 35 3-(3,5-dimethyl-4-ethoxycarbonylpyrrol-2-yl)-2-indolinone;

Benzoic acid, 2-[[[1-ethyl-2,3-dihydro-2-oxo-3-(1H-pyrrol-2-ylmethylene)-1H-indol-5-yl]oxy]methyl]-;  
3-[(1-methyl-5-nitro-imidazol-2-yl)methylene]-2-indolinone;

5 1-methyl-3-[(1-methyl-5-nitro-imidazol-2-yl)methylene]-2-indolinone;


1-ethyl-3-[(1-methyl-5-nitro-imidazol-2-yl)methylene]-2-indolinone;

3-(thien-2-ylmethylene)-2-indolinone;

10 1H-Indole-7-acetic acid, 3-[(2-butyl-1H-imidazol-4-yl)methylene]-2,3-dihydro-2-oxo-, ethyl ester;  
and

1H-Indole-7-acetic acid, 3-[[2-butyl-1-[(1,1-dimethylethoxy)carbonyl]-1H-imidazol-4-yl)methylene]-2,3-  
15 dihydro-2-oxo-, ethyl ester.

9. A compound of the formula:



and pharmaceutically acceptable salts thereof, wherein:

R<sub>1</sub> is H or alkyl;

R<sub>2</sub> is O or S;

30 R<sub>3</sub> is hydrogen;

R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>7</sub> are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH,  
35 CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';

R<sub>3</sub>', R<sub>4</sub>', R<sub>5</sub>', and R<sub>6</sub>' are each independently selected from the group consisting of hydrogen, alkyl, alkoxy,

aryl, aryloxy, alkaryl, alkaryloxy, halogen,  
trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH,  
CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';

n is 0-3;

5 R is H, alkyl or aryl; and


R' is H, alkyl or aryl.

10. The compound of claim 9 wherein:

R<sub>3</sub>' is selected from the group consisting of methyl,  
10 ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-  
butyl, halogen, aryl and OR, where R is H, alkyl or aryl;  
and

R<sub>5</sub>' is selected from the group consisting of  
hydrogen, methyl, ethyl, n-propyl, iso-propyl, n-butyl,  
15 iso-butyl, tert-butyl, halogen, aryl and OR, where R is  
H, alkyl or aryl.

11. A compound of the formula:



and pharmaceutically acceptable salts thereof, wherein:

R<sub>1</sub> is H or alkyl;

30 R<sub>2</sub> is O or S;

R<sub>3</sub> is hydrogen;

R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>7</sub> are each independently selected  
from the group consisting of hydrogen, alkyl, alkoxy,  
aryl, aryloxy, alkaryl, alkaryloxy, halogen,  
35 trihalomethyl, S(O)R, SO<sub>2</sub>NRR', SO<sub>2</sub>R, SR, NO<sub>2</sub>, NRR', OH,  
CN, C(O)R, OC(O)R, NHC(O)R, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R, and CONRR';

$R_2'$ ,  $R_3'$ ,  $R_5'$ , and  $R_6'$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $5\text{ CN}$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$n$  is 0-3;

$Z$  is Br, Cl, F, I, methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-butyl, or tert-butyl;

$R$  is H, alkyl or aryl; and

10  $R'$  is H, alkyl or aryl.

12. The compound of claim 11 wherein:

$R_3'$  is selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where  $R$  is H, alkyl or aryl; and


$R_5'$  is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, halogen, aryl and OR, where  $R$  is 20 H, alkyl or aryl.

13. A pharmaceutical composition comprising a pharmaceutically acceptable carrier or excipient and a compound according to Claim 1, 2, 9, 10, 11 or 12.

25 14. A method for treating diseases related to unregulated tyrosine kinase signal transduction, the method comprising the step of administering to a subject in need thereof a therapeutically effective amount of a 30 compound selected from the group consisting of:

(i) a compound of the formula:

5



10 and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$R_3'$ ,  $R_5'$ , and  $R_6'$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$n$  is 0-3;


25  $X$  is Br, Cl, F or I;

$R$  is H, alkyl or aryl; and

$R'$  is H, alkyl or aryl;

(ii) a compound of the formula:

30



35

and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

5  $R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

10  $R_2'$ ,  $R_3'$ ,  $R_5'$ , and  $R_6'$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

15  $n$  is 0-3;


$R$  is H, alkyl or aryl;

$R'$  H, alkyl or aryl; and

$R_a$  and  $R_b$  are each independently selected from the group consisting of H, alkyl and  $C(O)R$ , or  $NR_aR_b$  taken  
20 together are a heterocyclic ring of from 3 to 8 atoms optionally substituted at one or more positions with hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ ,  
25 and  $CONRR'$ ;

(iii) a compound of the formula:

30



35

and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

A is a five membered heteroaryl ring selected from the group consisting of thiophene, pyrrole, pyrazole, imidazole, 1,2,3-triazole, 1,2,4-triazole, oxazole, isoxazole, thiazole, isothiazole, 2-sulfonylfuran, 4-alkylfuran, 1,2,3-oxadiazole, 1,2,4-oxadiazole, 1,2,5-oxadiazole, 1,3,4-oxadiazole, 1,2,3,4-oxatriazole, 1,2,3,5-oxatriazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,3,4-thiatriazole, 1,2,3,5-thiatriazole, and tetrazole, optionally substituted at one or more positions with alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$  or  $CONRR'$ ;


$n$  is 0-3;

$R$  is H, alkyl or aryl; and

$R'$  is H, alkyl or aryl;

(iv) a compound of the formula:

30



35

and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$R_1'$ ,  $R_4'$ ,  $R_5'$ , and  $R_6'$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$n$  is 0-3;


$R$  is H, alkyl or aryl; and

$R'$  is H, alkyl or aryl; and

(v) a compound of the formula:

20

25



and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

30  $R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;



$R_2'$ ,  $R_3'$ ,  $R_4'$ , and  $R_5'$  are each independently selected from the group consisting of hydrogen, alkyl, alkyl, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$n$  is 0-3;

$Z$  is Br, Cl, F, I, methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-butyl, or tert-butyl;

$R$  is H, alkyl or aryl; and

10  $R'$  is H, alkyl or aryl.

15. The method of claim 14 wherein said disease is selected from the group consisting of cancer, blood vessel proliferative disorders, fibrotic disorders, 15 mesangial cell proliferative disorders and metabolic diseases.

16. The method of claim 14 wherein the blood vessel proliferative disorder is selected from the group 20 consisting of arthritis and restenosis.

17. The method of claim 14 wherein the fibrotic disorder is selected from the group consisting of hepatic cirrhosis and atherosclerosis.

25


18. The method of claim 14 wherein the mesangial cell proliferative disorder is selected from the group consisting of glomerulonephritis, diabetic nephropathy, malignant nephrosclerosis, thrombotic microangiopathy 30 syndromes, transplant rejection and glomerulopathies.

19. The method of claim 14 wherein the metabolic disorder is selected from the group consisting of psoriasis, diabetes mellitus, wound healing, inflammation 35 and neurodegenerative diseases.

20. A method for regulating tyrosine kinase signal transduction comprising administering to a subject a therapeutically effective amount of a compound selected from the group consisting of:

5 (i) a compound of the formula:

10



15 and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$R_3'$ ,  $R_5'$ , and  $R_6'$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$n$  is 0-3;


30  $X$  is Br, Cl, F or I;

$R$  is H, alkyl or aryl; and

$R'$  is H, alkyl or aryl;


35

(ii) a compound of the formula:



- 5
- 10 and pharmaceutically acceptable salts thereof, wherein:
- $R_1$  is H or alkyl;
- $R_2$  is O or S;
- $R_3$  is hydrogen;
- 15  $R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;
- 20  $R_2'$ ,  $R_3'$ ,  $R_4'$ , and  $R_5'$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;
- 25  $n$  is 0-3;
- $R$  is H, alkyl or aryl;
- $R'$  H, alkyl or aryl; and
- $Ra$  and  $Rb$  are each independently selected from the group consisting of H, alkyl and  $C(O)R$ , or  $NRaRb$  taken
- 30 together are a heterocyclic ring of from 3 to 8 atoms optionally substituted at one or more positions with hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ ,  $CONRR'$ ;
- 35

(iii) a compound of the formula:



10

and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

15

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

20

$A$  is a five membered heteroaryl ring selected from the group consisting of thiophene, pyrrole, pyrazole, imidazole, 1,2,3-triazole, 1,2,4-triazole, oxazole, isoxazole, thiazole, isothiazole, 2-sulfonylfuran, 4-alkylfuran, 1,2,3-oxadiazole, 1,2,4-oxadiazole, 1,2,5-oxadiazole, 1,3,4-oxadiazole, 1,2,3,4-oxatriazole, 1,2,3,5-oxatriazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,3,4-thiatriazole, 1,2,3,5-thiatriazole, and tetrazole, optionally substituted at one or more

30

positions with alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$  or  $CONRR'$ ;


$n$  is 0-3;

35

$R$  is H, alkyl or aryl; and

$R'$  is H, alkyl or aryl;

(iv) a compound of the formula:



and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;


$R_3'$ ,  $R_4'$ ,  $R_5'$ , and  $R_6'$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

$n$  is 0-3;

$R$  is H, alkyl or aryl; and

$R'$  is H, alkyl or aryl; and

(v) a compound of the formula:



and pharmaceutically acceptable salts thereof, wherein:

$R_1$  is H or alkyl;

$R_2$  is O or S;

$R_3$  is hydrogen;

5  $R_4$ ,  $R_5$ ,  $R_6$ , and  $R_7$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

10  $R_2'$ ,  $R_3'$ ,  $R_5'$ , and  $R_6'$  are each independently selected from the group consisting of hydrogen, alkyl, alkoxy, aryl, aryloxy, alkaryl, alkaryloxy, halogen, trihalomethyl,  $S(O)R$ ,  $SO_2NRR'$ ,  $SO_2R$ ,  $SR$ ,  $NO_2$ ,  $NRR'$ ,  $OH$ ,  $CN$ ,  $C(O)R$ ,  $OC(O)R$ ,  $NHC(O)R$ ,  $(CH_2)_nCO_2R$ , and  $CONRR'$ ;

15  $n$  is 0-3;

$Z$  is Br, Cl, F, I, methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-butyl, or tert-butyl;

$R$  is H, alkyl or aryl; and

$R'$  is H, alkyl or aryl.

20

25

30

35

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US96/08903

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/40; C07C 209/34

US CL :514/418; 548/486

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/418; 548/486

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,206,261 A (KAWAGUCHI ET AL.) 27 April 1993 (27.04.93), column 1, line 20 to column 2, line 53.	1-20
X	WO 92/07830 A (PFIZER, INC.) 14 May 1992 (14.05.92), page 1, line 31 to page 3, line 20.	1-20
X	EP 0,580,502 A1 (ADIR ET COMPAGNIE) 26 January 1992 (26.01.92), page 1, abstract.	1-20

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T	later documents published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	A*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

21 AUGUST 1996

Date of mailing of the international search report

03 SEP 1996

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RICHARD L. RAYMOND

Telephone No. (703) 308-1235

Form PCT/ISA/210 (second sheet) (July 1992)\*





conformational change in the nucleotide-binding loop. This structural information will facilitate the design of new inhibitors for use in the treatment of cancer and other diseases in which cell signaling by tyrosine


kinases plays a crucial role in disease pathogenesis.

IT 186611-14-3, SU 5402

RL: BPR (Biological process); BIOL (Biological study); PROC (Process) (inhibitor; structure of tyrosine kinase domain of fibroblast growth factor receptor in complex with inhibitors)

RN 186611-14-3 CAPLUS

CN 1H-Pyrrole-3-propanoic acid, 5-[(1,2-dihydro-2-oxo-3H-indol-3-ylidene)methyl]-4-methyl- (9CI) (CA INDEX NAME)



L4 ANSWER 14 OF 19 CAPLUS COPYRIGHT 1999 ACS

ACCESSION NUMBER: 1997:140244 CAPLUS

DOCUMENT NUMBER: 126:139901

TITLE: Indolinone compounds capable of modulating tyrosine kinase signal transduction

INVENTOR(S): Tang, Peng Cho; Sun, Li; McMahon, Gerald

PATENT ASSIGNEE(S): Sugen, Inc., USA

SOURCE: PCT Int. Appl., 133 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 6

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9640116	A1	19961219	WO 1996-US8903	19960605
W:	AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IL, IS, JP, KG, KP, KR, KZ, LK, LR, LS, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, UZ, VN, AM, AZ, BY			
RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
US 5880141	A	19990309	US 1995-485323	19950607
CA 2192797	AA	19961219	CA 1996-2192797	19960605
AU 9660441	A1	19961230	AU 1996-60441	19960605
AU 706597	B2	19990617		
EP 769947	A1	19970502	EP 1996-918093	19960605
R:	AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE			



BR 9606410 A 19971230 BR 1996-6410 19960605  
 JP 10504323 T2 19980428 JP 1996-501363 19960605  
 EP 934931 A2 19990811 EP 1999-103667 19960605  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI  
 NO 9605377 A 19970212 NO 1996-5377 19961213  
 PRIORITY APPLN. INFO.: US 1995-485323 19950607  
 EP 1996-918093 19960605  
 WO 1996-US8903 19960605

OTHER SOURCE(S): MARPAT 126:139901

AB The present invention relates to org. mols. capable of modulating tyrosine

kinase signal transduction in order to regulate, modulate and/or inhibit abnormal cell proliferation. Representatives of the 5 different classes of compds. described are SU 4932 [3-(2-chloro-4-hydroxybenzylidenyl)-2-indolinone], SU 4312 [3-(4-dimethylaminobenzylidenyl)-2-indolinone], SU 5416 [3-[(2,3-dimethylpyrrol-5-yl)methylene]-2-indolinone], SU 5204 [3-(2-ethoxybenzylidenyl)-2-indolinone], and SU 4942 [3-(4-bromobenzylidenyl)-2-indolinone]. Diseases which these compds. and their pharmaceutically acceptable prepsns. may be effective against include arthritis, hepatic cirrhosis, diabetic nephropathy and psoriasis.


IT 15966-93-5P, SU 5408 91E22-51-4P, SU 4314  
 186610-93-5P, SU 5404 186610-94-6P, SU 5406  
 186610-95-7P, SU 5416 186611-14-3P, SU 5402  
 186611-15-4P, SU 5403 186611-16-5P, SU 5405  
 186611-17-6P, SU 5407 186611-29-0P, SU 5453  
 186611-30-3P, SU 5454 186611-31-4P, SU 5455  
 186611-32-5P, SU 5456 186611-33-6P, SU 5459  
 186611-34-7P, SU 5460 186611-35-8P, SU 5461  
 186611-36-9P, SU 5462 186611-37-0P, SU 5463  
 186611-38-1P, SU 5464 186611-39-2P, SU 5465  
 186611-48-3P, SU 5477 186611-49-4P, SU 5478  
 186611-50-7P, SU 5479 186611-53-0P, SU 5612  
 186611-54-1P, SU 5613 186611-56-3P, SU 5614  
 186611-65-4P, SU 5624 186611-66-5P, SU 5625  
 186611-67-6P, SU 5626

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(prepn. of indolinones capable of modulating tyrosine kinase signal transduction)

RN 15966-93-5 CAPLUS


CN 1H-Pyrrole-3-carboxylic acid, 5-[(1,2-dihydro-2-oxo-3H-indol-3-ylidene)methyl]-2,4-dimethyl-, ethyl ester (9CI) (CA INDEX NAME)









RN 91822-51-4 CAPLUS  
 CN 2H-Indol-2-one, 1,3-dihydro-3-((3-methyl-1H-pyrrol-2-yl)methylene)- (9CI) (CA INDEX NAME)



RN 186610-93-5 CAPLUS  
 CN 2H-Indol-2-one, 1,3-dihydro-3-((1H-pyrrol-2-yl)methylene)- (9CI) (CA INDEX NAME)




RN 186610-94-6 CAPLUS  
 CN 2H-Indol-2-one, 3-((3,4-dimethyl-1H-pyrrol-2-yl)methylene)-1,3-dihydro- (9CI) (CA INDEX NAME)




RN 186610-95-7 CAPLUS  
 CN 2H-Indol-2-one, 3-((4,5-dimethyl-1H-pyrrol-2-yl)methylene)-1,3-dihydro- (9CI) (CA INDEX NAME)






RN 186611-14-3 CAPLUS

CN 1H-Pyrrole-3-propanoic acid, 5-[(1,2-dihydro-2-oxo-3H-indol-3-ylidene)methyl]-4-methyl- (9CI) (CA INDEX NAME)




RN 186611-15-4 CAPLUS

CN 2H-Indol-2-one, 3-[(3,4-dibromo-5-methyl-1H-pyrrol-2-yl)methylene]-1,3-dihydro- (9CI) (CA INDEX NAME)



RN 186611-16-5 CAPLUS

CN 1H-Pyrrole-2-carboxaldehyde, 5-[(1,2-dihydro-2-oxo-3H-indol-3-ylidene)methyl]-3,4-dimethyl- (9CI) (CA INDEX NAME)




RN 186611-17-6 CAPLUS

CN 1H-Pyrrole-3-propanoic acid, 2-[(1,2-dihydro-2-oxo-3H-indol-3-ylidene)methyl]-4-methyl-, methyl ester (9CI) (CA INDEX NAME)









RN 186611-29-0 CAPLUS

CN 2H-Indol-2-one, 3-[(3-ethyl-4,5-dimethyl-1H-pyrrol-2-yl)methylene]-1,3-dihydro- (9CI) (CA INDEX NAME)




RN 186611-30-3 CAPLUS

CN 1H-Pyrrole-3-propanoic acid, 5-[(1,2-dihydro-2-oxo-3H-indol-3-ylidene)methyl]-2-(ethoxycarbonyl)-4-(2-ethoxy-2-oxoethyl)-, ethyl ester (9CI) (CA INDEX NAME)




RN 186611-31-4 CAPLUS

CN 1H-Pyrrole-2-carboxylic acid, 5-[(1,2-dihydro-2-oxo-3H-indol-3-ylidene)methyl]-4-ethyl-3-methyl- (9CI) (CA INDEX NAME)







RN 186611-32-5 CAPLUS  
 CN 2H-Indol-2-one, 3-[(3,5-diiodo-4-methyl-1H-pyrrol-2-yl)methylene]-1,3-dihydro- (9CI) (CA INDEX NAME)



RN 186611-33-6 CAPLUS  
 CN 1H-Pyrrole-3-acetic acid, 2-chloro-5-[(1,2-dihydro-2-oxo-3H-indol-3-ylidene)methyl]-4-(methoxycarbonyl)-, methyl ester (9CI) (CA INDEX NAME)




RN 186611-34-7 CAPLUS  
 CN 1H-Pyrrole-2-carboxylic acid, 4-acetyl-5-[(1,2-dihydro-2-oxo-3H-indol-3-ylidene)methyl]-3-methyl-, ethyl ester (9CI) (CA INDEX NAME)




RN 186611-35-8 CAPLUS  
 CN 2H-Indol-2-one, 3-[[1-(3,5-dichlorophenyl)-1H-pyrrol-2-yl]methylene]-1,3-dihydro- (9CI) (CA INDEX NAME)






RN 186611-36-9 CAPLUS

CN 2H-Indol-2-one, 3-[[1-(4-chlorophenyl)-1H-pyrrol-2-yl]methylene]-1,3-dihydro- (9CI) (CA INDEX NAME)




RN 186611-37-0 CAPLUS

CN 1H-Pyrrole-3-carboxylic acid, 5-[(1,2-dihydro-2-oxo-3H-indol-3-ylidene)methyl]-4-methyl-, ethyl ester (9CI) (CA INDEX NAME)



RN 186611-38-1 CAPLUS


CN 2H-Indol-2-one, 1,3-dihydro-3-[(1-methyl-1H-pyrrol-2-yl)methylene]- (9CI) (CA INDEX NAME)






RN 186611-39-2 CAPLUS

CN 1H-Pyrrole-3-propanoic acid, 2-[(1,2-dihydro-2-oxo-3H-indol-3-ylidene)methyl]-5-(ethoxycarbonyl)-4-(2-ethoxy-2-oxoethyl)-, ethyl ester (9CI) (CA INDEX NAME)




RN 186611-48-3 CAPLUS

CN 2H-Indol-2-one, 3-[(4-ethyl-3,5-dimethyl-1H-pyrrol-2-yl)methylene]-1,3-dihydro- (9CI) (CA INDEX NAME)




RN 186611-49-4 CAPLUS

CN 2H-Indol-2-one, 3-[(5-ethyl-1H-pyrrol-2-yl)methylene]-1,3-dihydro- (9CI) (CA INDEX NAME)




RN 186611-50-7 CAPLUS

CN 2H-Indol-2-one, 3-[[3,5-dimethyl-4-(1-methylethenyl)-1H-pyrrol-2-yl)methylene]-1,3-dihydro- (9CI) (CA INDEX NAME)











RN 186611-53-0 CAPLUS

CN 2H-Indol-2-one, 5-chloro-1,3-dihydro-3-(1H-pyrrol-2-ylmethylene)- (9CI)  
(CA INDEX NAME)


RN 186611-54-1 CAPLUS

CN 2H-Indol-2-one, 5-chloro-1,3-dihydro-3-[(3-methyl-1H-pyrrol-2-yl)methylene]- (9CI) (CA INDEX NAME)



RN 186611-56-3 CAPLUS


CN 2H-Indol-2-one, 5-chloro-3-[(3,5-dimethyl-1H-pyrrol-2-yl)methylene]-1,3-dihydro- (9CI) (CA INDEX NAME)



RN 186611-65-4 CAPLUS


CN 2H-Indol-2-one, 1,3-dihydro-5-nitro-3-(1H-pyrrol-2-ylmethylene)- (9CI)  
(CA INDEX NAME)






RN 186611-66-5 CAPLUS

CN 2H-Indol-2-one,  
1,3-dihydro-3-[(3-methyl-1H-pyrrol-2-yl)methylene]-5-nitro-  
(9CI) (CA INDEX NAME)



RN 186611-67-6 CAPLUS

CN 2H-Indol-2-one, 3-[(3,5-dimethyl-1H-pyrrol-2-yl)methylene]-1,3-dihydro-5-nitro- (9CI) (CA INDEX NAME)



L4 ANSWER 15 OF 19 CAPLUS COPYRIGHT 1999 ACS

ACCESSION NUMBER: 1992:490138 CAPLUS

DOCUMENT NUMBER: 117:90138

TITLE: Preparation of 3-(arylmethylene)oxindoles and analogs  
as gastrin releasing peptide antagonists

INVENTOR(S): Nakanishi, Susumu

PATENT ASSIGNEE(S): Pfizer Inc., USA

SOURCE: PCT Int. Appl., 17 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9207830	A2	19920514	WO 1991-US4978	19910718

